


CAUTION: Protect children, domestic animals, and wild life from the bait. If swallowed by humans, domestic animals or wild life it may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of EshydroxyCoumarin. Bait should be stored in areas not accessible to children and pets, children and domestic animals or in tamper proof bait boxes.

Rat Rid WATER SOLUBLE BAIT

KEEP AWAY FROM DOGS, CATS AND OTHER DOMESTIC ANIMALS AND WILDLIFE. KEEP AWAY FROM AROMATIC SUBSTANCES. Do not use in food areas. Destroy when empty.



RAT RID has been field tested in areas densely infested with rats, and which offered a liberal selection of foods. *RAT RID* is *ATTRACTIVE, ACCEPTABLE, EFFECTIVE*. *ATTRACTIVENESS*, field tests have shown that *RAT RID* is highly acceptable.

ACCEPTABILITY, field tests have shown that once rats begin to feed on *RAT RID* it will produce a lethal dose.

EFFECTIVENESS, field tests have shown that *RAT RID* will kill rats and mice when used as directed.

RESULTS, Long Term Control by continuous baiting with *RAT RID*.

E.P.A. Reg. No. 1266-116
ACTIVE INGREDIENT:
Warfarin (3,3'-Aceton, Benzylidene-
hydroxy-coumarin) 0.025
INERT INGREDIENTS 99.975

CAUTION
KEEP OUT OF REACH OF CHILDREN
See back panel for additional directions

ACCEPTED
May 24 1972
U.S. DEPT. OF AGRICULTURE
FEDERAL AGENCY FOR PROTECTION OF THE ENVIRONMENT
WASHINGTON, D.C. 20250

1266-116

DIRECTIONS

RATS: Use 2 *RAT RID* bait boxes per room. Bait boxes with rats feed, water, and or travel. Placement should be in areas where rats are likely to feed. Check bait stations daily to assure an uninterrupted supply of bait for not less than 1 day. Continue until all signs of feeding have stopped.

MICE: Place 1 *RAT RID* bait box in a shallow, dry container next to burrows, in runways, or other places where mice are likely to feed. It is recommended that bait be placed at intervals of 8 to 12 feet. Check bait stations daily to assure an uninterrupted supply of bait for not less than 15 days. Continue until all signs of feeding have stopped.

CONSTANT CONTROL: Replace depleted bait. Stake down or moidy baits should be replaced with fresh bait promptly. Bait should be placed in locations not accessible to children, pets, and domestic animals. For tamper proof bait boxes, **CONTROL: Controls Norway Rat, Roof Rat and House Mice.**