

NOV 27 1983

300 / 104594  
17 / 2

Malter International Corporation  
P.O. Box 6099  
New Orleans, LA 70174

Gentlemen:

Subject: SEC-T-CIDE  
EPA Registration No. 1266-58  
Your Application Dated August 26, 1983

Your proposed amendment to update the labeling for this product is acceptable. A stamped copy of the draft labeling is enclosed for your records. Submit five (5) copies of the finished printed labeling, corrected in accordance with the comments listed below, before the product is released for shipment under this amended labeling.

- (1) Add the following statement to the precautionary labeling: "Remove birds and pets and cover or remove fish aquariums before spraying."
- (2) Change the First Aid section to read:

If swallowed, do not induce vomiting. Call a physician. For eye contact, flush with plenty of water and see a physician.

Sincerely yours,

Timothy A. Gardner  
Product Manager (17)  
Insecticide-Rodenticide Branch  
Registration Division (48-767)

Enclosure

RD: ~~SAIT/WANG~~ DCR-12111; WANG-1268C; vgr; kaven: 479-2013: 11/25/83

CONCURRENCES							
SYMBOL	TS 767						
DATE	11/25/83						

CENTER PANEL

NOV 28 1983

For Professional and  
Industrial Use Only

MALTER  
INTERNATIONAL

1266-58

S E C - T - C I D E

\*Non-Residual      \*Kills By Contact

\*Indoor And Outdoor Use

E.P.A.Reg. No. 1266-58-2A  
E.P.A.Est. No. 1266-LA-1

ACTIVE INGREDIENTS:

Petroleum Distillate . . . . .	98.35%
*Technical Piperonyl Butoxide . . . . .	0.90%
Pyrethrins . . . . .	0.18%

INERT INGREDIENTS: . . . . . 0.57%

\*Equivalent to 0.72% (butylcarbityl) (6-propylpiperonyl)  
either and 0.18% related compounds.

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

(See left panel for additional precautionary and  
practical treatment statements)

MALTER INTERNATIONAL CORPORATION 1973  
International Headquarters:  
P. O. Box 6099, New Orleans, Louisiana 70174 U.S.A.  
Telephone: 504/362-3232 - Telex No.:58-4184

2

LEFT PANEL

HAZARDS TO HUMANS & DOMESTIC  
ANIMALS

**CAUTION:** Harmful if swallowed. Avoid contamination of feed and foodstuffs. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when the facility is not in operation. Do not spray animals or pets.

**FIRST AID:** If swallowed, do not induce vomiting and call physician. For eye contact, flush with water for 15 minutes and see physician.

**PHYSICAL AND CHEMICAL HAZARDS:** Do not use, pour, spill, or store near heat or open flame.

**ENVIRONMENTAL HAZARDS:** This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of spray equipment or disposal of wastes.

**DIRECTIONS FOR USE:**

**GENERAL CLASSIFICATION:**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**SEC-T-CIDE SPACE SPRAY** is an excellent insecticide for use as a vapor, fog or mist spray in mechanical generators and electric vaporizers, such as Malter's Portable Electro Fogger. Application should be sufficient to fill the enclosure with fog, mist or vapor. When using spray in food processing areas, use only when the plant is not in operation.

Shut off ventilating systems and air conditioning units before spraying. Cover food, food products, utensils, etc., to prevent contamination from falling insects. Treat infested areas thoroughly and whenever possible. Keep doors and windows closed for 10 minutes after spraying. Do not remain in treated areas. Ventilate the area after treatment is completed. Then sweep up and dispose of fallen insects. Spray as often as necessary to control these insects.

## DIRECTIONS (CONTINUED)

**Flying Insects:** For control of flies, mosquitoes, gnats, wasps, hornets, and flies: direct the spray towards the upper portions of the room and saturate the atmosphere with the mist, using 1 ounce to each 3,000 cubic feet of space. Rapidity of initial kill may be accelerated by increasing dosage; however, not more than 1 gallon should be atomized in each 40,000 cubic feet.

**Crawling Insects:** To control roaches, ants, silverfish, crickets, spiders, spray infested areas, thoroughly and direct the liquid behind baseboards, moldings, cabinets, shelves, sinks, door and window frames and pipes, toward drains and dark corners and into cracks and similar hiding places. For best results contact insects directly whenever possible. Repeat treatment when necessary.

**In controlling fleas and brown dog ticks in houses:** Direct the application toward the lower wall portions, baseboards, rugs, carpets and floors of infested areas. Animal quarters, kennels and bedding should be sprayed.

For most effective control of clothes moths and carpet beetles apply spray liberally in empty chests, closets, bureaus and storages; direct the liquid into cracks, joints and crevices. Infested articles must be treaded thoroughly by holding the spray nozzle at least 3 feet away from the fabric and applying the liquid lightly on the entire article, paying particular attention to the seams and folds. Treat both sides of rugs, carpets and pads, and all exposed and accessible surfaces of overstuffed furniture. Repeat treatment monthly.

**To control bedbugs effectively:** Apply liberally to top, ends, sides and bottom of mattresses, spraying thoroughly all tufts, seams, and folds. Spray springs and frame, directing the liquid into all cracks and lock joints. Treat walls, baseboards, moldings, floor, closets and bedroom furniture, with particular attention to protected bedbug hiding and breeding places.

(over)

(Con't)

4

For home and institutional pantries and food storage infested with GRANARY WEEVIL, CADELLES, RICE WEEVIL, MEAL WORMS, SPIDER BEETLES, FLOUR BEETLES, RUST RED FLOUR BEETLE, SAWTOOTHED GRAIN BEETLE, MEAL MOTHS, SKIPPERS, GRAIN AND CHEESE MITES: Foods should be removed or covered during treatment. Spray thoroughly all surfaces of shelves, bins, cupboards, cabinets, pantries, lockers, and similar storages, with particular attention to cracks and other hiding places. Repeat treatment as needed to prevent infestations.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Reseal container and offer for reconditioning or triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill or bury in a safe place. Consult Federal, state or local disposal authorities for approved alternative procedures.