

DIRECTIONS:

Lawns and Ornamental Turf — To control dandelions, chickweed, plantain, Japanese clover, oxalis, wild onions and similar weeds, dilute 2 quarts DI-PHEN with 100 gallons of water to treat one acre. Water thoroughly before application. Wet foliage thoroughly with solution. Do not apply during heat of day. If grass begins to yellow, water thoroughly and fertilize. Do not apply to grass only 3 or 4 months old. Temporary burning or yellowing may occur to Bent and Fescue grasses. Do not use on Clover, Dichondra, Ice Plants or Lippia lawns. For hard to kill weeds such as Oxalis, repeat treatment may be necessary. Wild Onions and similar weeds — Treat while young and before bulbs are formed, usually in late fall or early spring.

NOTE: CARPETGRASS AND ST. AUGUSTINE GRASS MAY BE INJURED. APPLY ONLY AS A SPOT TREATMENT. DO NOT USE ON ICE PLANTS.

For Docks, Thistles, Morning Glory in Fence rows and rights of way — Mix 1 gallon to 100 gallons of water to treat 1 acre. Wet plants thoroughly and repeat when necessary.

For woody plants such as Poison Oak, Brambles, Poison Ivy, Sumac, Honeysuckle: Mix 8 ounces per gallon of water and drench thoroughly while plants are in full leaf. Repeat application in four weeks if necessary.

To Kill Stumps — Mix 1 pint in one quart of Kerosene or Diesel Oil. Spray all of the stump area thoroughly.

NOTICE

Keep from freezing. If this product is stored in cold or freezing weather it may separate or thicken. This will not harm the performance. Before using, allow to warm to room temperature and stir thoroughly.



DI-PHEN

BROADLEAF WEED & WOODY BRUSH KILLER

• SELECTIVE • LOW VOLATILE

ACTIVE INGREDIENTS

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

MALTER INTERNATIONAL CORPORATION

1000 N. 10TH AVE. HEALD JUNCTION, P.O. BOX 5099, NEW ORLEANS, LA 70112

CAUTION

Avoid contact with skin, eyes or clothing. Harmful if swallowed. In case of contact flush with plenty of water, wash with soap and water. Avoid breathing spray mist. KEEP OUT OF REACH OF CHILDREN.

DO NOT allow solution or spray drift to contact desirable plants. Among crops highly susceptible to DI-PHEN injury are cotton, tobacco, blackeye peas, beans, tomatoes, melons, other vegetables, grapes, fruit trees and some ornamentals. (Coarse sprays are less likely to drift than are fine mist sprays.)

It is difficult to completely clean DI-PHEN from sprayers or other equipment, therefore do not use such sprayers and other equipment for purposes where even trace amounts of this chemical may cause injury.

Do not reuse shipping containers; destroy by crushing and burying. Avoid contamination of water intended for irrigation and domestic use. Do not transport with or store near seeds, fertilizers, insecticides or fungicides.

Vapors from this product may injure susceptible plants in the vicinity.

Do not use, pour, spill or store near heat or open flame. Consult local Experiment Station or Extension Service, Weed Specialists for specific recommendations as to dosage and time of application and for control of other weeds and woody plants. Note: Keep spray tank agitated.

Apply only to areas that will not be grazed after treatment by milking animals or animals being finished for slaughter.


Copyright MCMLXXII. Printed in U.S.A.

CENTER PANEL

D I - P H E N

BROADLEAF WEED
AND WOODY
BRUSH KILLER

Selective, One-Step Weed Control




Warning! KEEP OUT OF REACH OF CHILDREN. Flammable. Contents under pressure.
See left panel for additional precautions. Net. Wt. 13 oz. (369 grams)

LEFT PANEL

WARNING! Flammable! Contents under pressure. Do not use near fire, sparks, or flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Use with adequate ventilation. Do not breathe mist. Avoid contact with skin, eyes or clothing. In case of contact, flush with plenty of water or wash with soap and water. Do not allow spray drift to contact desirable plants. KEEP OUT OF REACH OF CHILDREN.

RIGHT PANEL

D I - P H E N
BROADLEAF WEED
AND WOODY
BRUSH KILLER



Selective, low volatile weed killer controls weeds, brush and woody plants. DI-PHEN is recommended to kill and control: Brambles, Chickweed, Dandelions, Docks, Honeysuckle, Japanese Clover, Morning Glory, Oxalis, Plantain, Poison Ivy, Poison Oak, Poison Sumac, Thistles and Wild Onion.

E.P.A. Reg. No. 1266-151
E.P.A. Est. No. 1266-LA-1

ACTIVE INGREDIENTS:

Petroleum Distillates..... 72.20%
Isooctyl Ester of 2,4-Dichlorophenoxy acetic Acid*.. 02.36%
Isooctyl Ester of Silvex 2(2,4,5-Trichlorophenoxy
propionic Acid**..... 01.05%
INERT INGREDIENTS: 24.39%
* Equivalent to 1.56% 2,4-D Acid
** Equivalent to 0.7% Silvex

D I R E C T I O N S :

DI-PHEN is to be used for localized or spot treatment of areas where broadleaf weeds or woody-brush weeds are not desirable. Do not use near desirable plants.

Shake well before using. Spray weeds thoroughly from a distance of about 2 ft. Repeat treatment if regrowth occurs.