

[1266-105]

CAUTION:

Harmful if swallowed. Keep out of reach of children. THIS INSECTICIDE IN THE CONCENTRATED FORM IS FLAMMABLE AND SHOULD NOT BE USED, POURED, SPILLED, OR STORED NEAR HEAT, SPARKS, OR AN OPEN FLAME. REMOVE PETS, BIRDS, AND COVER FISH AQUARIUMS BEFORE SPRAYING.

DIRECTIONS:

For Food and Beverage plants and feed storage areas for control of mosquitoes, wasps, flies, flying moths, ants, roaches, spiders, and silverfish. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product as a space spray in these areas, apply only when plant is not in operation and remove or cover all foods.

For space spraying of Buildings and Enclosures for mosquitoes, wasps, flies, and flying moths: Dilute 1 gallon of DAIRY-CIDE with 5 gallons of water. Close all doors and windows. Thoroughly spray all parts of the room, particularly toward the ceiling, hitting as many insects as possible and continue spraying until the room is filled with a fine mist. Leave room closed for 15 to 20 minutes. Do not remain in treated areas and ventilate area after treatment is completed.

Crawling insects (ants, roaches, spiders, and silverfish); Dilute 2 parts by volume of water. Use a sprayer developing a coarse, wet spray. Treat cracks and crevices in woodwork, dark places behind pipes, and all places where the insects infest, hitting as many as possible.



ACCEPTED
29 JUL 1974
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 1266-105 SUBJECT TO ATTACHED COMMENTS

DAIRY-CIDE

HIGHLY CONCENTRATED NON-RESIDUAL WATER EMULSIFIABLE INSECTICIDE

E.P.A. Reg. No 1266-105-AA
E.P.A. Est. No. 1266-LA-1

ACTIVE INGREDIENTS:94%
Petroleum Distillates, Methylated Naphthalene, Technical Piperonyl Butoxide*, Pyrethrins
*Equivalent to (Butylcarbityl) (6-Propylpiperonyl)
Ether and related compounds.
INERT INGREDIENTS:6%
Dodecylphenyl Polyethylene Glycol Ether

CAUTION: KEEP OUT OF REACH OF CHILDREN.
See left panel for additional cautions.

© Copyright 1974 by:
MALTER INTERNATIONAL CORPORATION
INTERNATIONAL HEADQUARTERS: P.O. Box 6099, New Orleans, La. 7-174
Telephone 504 / 362-3232 - Telex No. 58 - 4134

DIRECTIONS (Continued):

Repeat treatment as required.
Important: Water diluted sprays; when used in the home, may cause stains to wallpaper, paints, and other surfaces.
MALTER'S DAIRY-CIDE when diluted with water, is non-flammable. When used as directed, it will kill the exposed stages of horn flies, stable flies, horse flies, deer flies, gnats, mosquitoes, wasps, house flies, cattle lice, spiders, silverfish, roaches, ants, and certain other insects associated with feed and food products.

For Spraying Cattle, Horn Flies, Mosquitoes, and Gnats: Dilute with 25 parts by volume of water and use two quarts of dilution per animal every five days or more if needed.

Stable Flies, Horse Flies, and Deer Flies: Dilute with 2 parts by volume of water and use one quart of dilution per animal on the legs, shoulders and undersides. Apply every two days or more often if necessary.

Cattle Lice: For single treatment to kill both young and adult lice, dilute 1 gallon of DAIRY-CIDE with 50 gallons of water and apply at a rate of 1 gallon or more per animal. Repeat in about 10 days or when necessary.

Do not reuse empty drum. Return to drum conditioner or destroy by perforating or crushing and burying in a safe place.

NOTICE: Keep from freezing. If this product is stored in cold or freezing weather it may separate or thicken. This will not harm the performance. Before using, allow to warm to room temperature and stir thoroughly.