

# Dewitt 25% Chlordane Emulsifiable Concentrate

## ACTIVE INGREDIENTS:

Technical chlordane 25%

Petroleum hydrocarbons 70%

## INERT INGREDIENTS:

Equivalent to 1% Octachloro-4,7-dimethyltricyclohexane and 10% related compounds

This is a concentrate insecticide for use as directed on label for use.

*For Roaches, Waterbugs, and Silverfish:* Mix 1 part with 12 1/2 parts water and apply thoroughly with coarse spray around shelves, bookcases, storage areas, infested cracks, and other hiding or resting places. Repeat as often as necessary to maintain effective control.

*For Ants:* Use same as for roaches, particularly on ant trails and crevices around door sills, window frames, openings around water pipes, heat ducts, and baseboards and where ants may enter the premises.

*For Brown Dog Ticks in Premises:* Use same as for roaches and particularly around sleeping quarters of pets and localized areas of floors and floor coverings. Place fresh bedding in animal quarters after treatment. Do not use directly on pets or other animals.

*For Carpet Beetles:* Use as for roaches on localized areas on floors, baseboards, and cracks under carpets where insects may be found. Repeat as needed.

*For Fleas, Ticks, Ants, and Chiggers:* In outdoor areas, mix 1 part with 12 1/2 parts of water, and apply the coarse wet spray in infested areas, except where food, feed and forage are grown. Wet thoroughly foliage of weeds and high grass.

*For Lawn and Ornamental Turf Insects Such as White Grubs, Japanese Beetle Larvae:* Mix 1 part concentrate with 10 parts of water and apply 1 gallon of the solution to 1,000 sq. ft. of lawn. Wash the insecticide into the soil with ample quantities of additional water. Keep children and pets off lawn until grass is completely dry. Birds feeding on treated areas may be harmed.

*For Wireworms and Mole Crickets, for Lawns:* Mix as above and apply 1/2 gallon to 1,000 sq. ft. washing in thoroughly.

*For Earwigs, and Sod Webworms:* First wet down the lawn thoroughly. Mix 1 part concentrate to 10 parts of water and apply this solution at a rate of 1 quart to 1,000 sq. ft. of lawn. Do not water lawn again for 2 to 3 days.

*For Corn Rootworms on Corn—Band Spraying:*—Use 2 quarts concentrate with sufficient water to cover 1 acre. Apply by spraying in bands of approximately 6 inches per row at planting time. For broadcast treatment, apply 2 gallons per acre with sufficient water to cover. Work into soil to depth of several inches.

*For Control of Flies and Maggots Around Outside Garbage and Refuse Dumps:* Mix 1 part concentrate with 24 parts water and apply as a coarse spray. Apply solution to point of run-off on garbage and refuse.

*For Tobacco Transplant Water:* Mix 1/2 to 1 pint (depending on degree of infestation of wireworms) to 50 gallons of water. Apply 200 to 300 gallons of this mix per acre.

Consult State Agricultural Extension Service for control of other crop insects.

*For Subterranean Termites:* Mix 1 part concentrate with 24 parts of water and thoroughly soak infested areas. If possible dig a trench 1 to 2 feet deep and 8 to 12 inches wide next to foundation wall; 30 inches in depth for buildings with deep foundations. Apply 2 gallons per 5 linear feet of trench for trenches up to 18 inches deep, and 4 gallons per 5 linear feet of trench for trenches over 18 inches deep. Apply at the rate of 2 gallons per 10 square feet in crawl spaces inside buildings and in critical areas, including cracks or breaks in masonry or concrete. Treat the soil as the trench is refilled. Treat piers under the house also. Inspect premises annually for reinfestation and repeat application as needed.

## CAUTION

### KEEP OUT OF THE REACH OF CHILDREN

USE ONLY AS DIRECTED. Apply only as specified on label. Harmful if swallowed.

Chlordane is toxic and may be absorbed through the skin. Wash thoroughly after use. Avoid inhaling spray and mists. Do not smoke while spraying. Avoid contamination of foodstuffs. Do not spray the undiluted material into or near fire or open flame.

Do not use on humans, household pets, or other animals. Do not use on dairy animals or other livestock or in dairy barns. Do not use on forage crops to be fed to dairy animals or animals being finished for slaughter. Toxic to fish and wildlife. Keep out of lakes, streams, and ponds. Rinse equipment and containers and dispose of waste by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning. (Keep out of smoke.)

*Antidote:* Call a physician immediately.

*Internal:* Emetic of mustard.

*External:* Wash with soap and warm water.

U.S.D.A. Reg. No. 1269-42

Net Contents: Gals.

Quality Control No.

# DE WITT CHEMICAL COMPANY

ATLANTA, GEORGIA, U. S. A.