



DE WITT

8D63

MANUFACTURERS OF
INDUSTRIAL MAINTENANCE
PRODUCTS

ACCEPTED
2-13-67
UNDER QUALITY CONTROL NO.
1269-48

CLEANER — DISINFECTANT — DEODORANT
For Hard Surfaces In and Around Chicken Hatcheries
GERMICIDAL — FUNGICIDAL — CLEANS AND DISINFECTS IN ONE OPERATION
KILLS STAPH AND TYPHOSA!

Comment

ACTIVE INGREDIENTS: 39.0%
Diethanolamine coconut soap; Isopropanol; Potassium ortho-benzyl-para-chlorophenate; Sodium ethylene diamine tetra-acetate; Trisodium phosphate; Potassium ortho-phenylphenate; Potassium 4-chloro-2-phenylphenate; Potassium 6-chloro-2-phenylphenate; Formaldehyde.

INERT INGREDIENTS: 61.0%
Water; Sodium tripolyphosphate; Perfume; Sodium nitrite.

Phenol Coefficients:
Staphylococcus aureus 15
Salmonella typhosa 18

Meets A.O.A.C. Use Dilution Test against Staphylococcus aureus and Salmonella choleraesuis at 1:128 dilution.

Non-flammable, Non-volatile.

DEWITT 8D63 is an effective combination of detergents, sanitizers, and deodorants for cleaning and sanitizing floors, walls, hatching and setting trays, incubators, egg trays, racks, and all other hard surfaces.

CAUTION

This is a concentrate. Dilute as directed above before use. Use hot water if available. Harmful if swallowed. Avoid contamination of feed or foodstuffs. May cause skin irritation. Avoid contact with skin, eyes, and clothing. In case of contact, flush with water; for eyes, get medical attention.

KEEP AWAY FROM CHILDREN.

DIRECTIONS FOR USE

1. To clean and disinfect hard surfaces such as those named above for chicken hatcheries, depending upon degree of soil, wash with solution containing from 1 to 3 ozs. DEWITT 8D63 per gallon of water. Allow solution to remain on surface 5 minutes. Rinse.
2. If surface has been cleaned, to use as a disinfectant *only* on above type surfaces, a dilution of 1/4 to 1 oz. per gallon of water is effective. Allow 5 minutes contact time.
3. Incubators may be sanitized, when not in use, by taking out egg trays, nursery trays, and other readily removable parts and scrubbing as directed in (1) above. Chick down, droppings, and shell fragments should be removed from the incubator with a brush or vacuum cleaner, and the interior sprayed with a 1 oz. per gallon solution of DW8D63. Allow to remain 5 minutes or more, and rinse with jet from water hose. Chick brooders and brooder rooms, when not in use, may be disinfected in a similar manner.
4. Chicken houses, enclosed runs, and all eating, drinking, and other utensils should be thoroughly cleaned and disinfected as in (1) above.

Controls the Environmental Spread of the Following Organisms:

- Pasteurella multocida (Fowl cholera)
- Salmonella gallinarum (Fowl typhoid)
- Salmonella pullorum (Pullorum disease— formerly called bacillary white diarrhea)
- S. typhosa
- Aspergillus fumigatus (Green mold)
- A. niger (Black mold)
- Staphylococcus aureus (Golden staph)

U.S.D.A. Reg. No. 1269-48

Net Content: GALS.

DE WITT CHEMICAL COMPANY

ATLANTA, GEORGIA, U. S. A.

ACCEPTED WITH COMMENTS