



DE WITT

Diazinon Emulsifiable Concentrate

FOR CONTROL OF COCKROACHES, FLIES AND OTHER HOUSEHOLD PESTS
FOR USE UNDER THE SUPERVISION OF TRAINED PEST CONTROL OPERATORS

ACTIVE INGREDIENTS:

O, O-Dimethyl O-(2, isopropyl-4-methyl-6-pyrimidinyl)	
phosphorothioate.....	12.5%
Aromatic petroleum derivative.....	82.5%
	95.0%
INERT INGREDIENT:	5.0%
	100.0%

DIRECTIONS:
Household Insects

Cockroaches, Water Bugs, Ants and Silverfish in Buildings: Prepare a 1% Diazinon spray by thoroughly mixing 5 ounces of DeWitt Diazinon with sufficient water to make one gallon of finished spray. Do not make up more spray than is to be used daily. Make spot application of this mixture to areas infested with these insects. Ant trails and places where ants enter the building are to be sprayed. Application should be made as a coarse spray with a paint brush to baseboards, areas around water pipes, cracks and crevices, surfaces behind and beneath cabinets, refrigerators, sinks, stoves and similar areas where these insects hide. Repeat at frequent intervals to prevent re-infestations.

Crack and Crevices in Buildings: Prepare a 1% Diazinon spray by thoroughly mixing 5 ounces of DeWitt Diazinon with sufficient water to make one gallon of finished spray. Do not make up more spray than is to be used daily. Thoroughly spray infested areas around baseboards, windows and door frames, wall cracks, sleeping quarters of household pets and localized areas of floors and floor coverings. Application should be repeated as infestations warrant and as re-infestations occur. Fresh bedding should be placed in animal quarters following treatment. Do not treat animals with this formulation. When used in dwellings, care should be taken to avoid deposits which could be frequently contacted by children. Care should also be taken to avoid spotting of wallpaper and fabrics. Do not contaminate food, food containers or cooking utensils.

Fly Control in Food Processing Plants: Use 10 ounces per gallon of water and apply as a residual spray to the walls and ceilings of food processing plants early in the season before canning or other processing begins. Repeat applications, if necessary, when plant is not in operation. All processing equipment and produce should be covered to avoid contamination. Spraying of unloading docks and other outside areas where flies congregate will aid in controlling flies.

CAUTION
KEEP OUT OF THE REACH OF CHILDREN.

May Be Fatal If Swallowed. May Be Absorbed Through Skin. Do not breathe spray mist. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed and feedstuffs. Do not use on livestock, household pets or humans. Do not permit children or pets to contact the treated surfaces until the spray has completely dried.

Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in service areas while food is exposed.

This product is toxic to fish and wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not use, pour, spill or store near heat or open flame.

Do not re-use empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies. Drums may be returned to drum reconditioner or disposed of as above.

Net Contents _____ Gallons Control No. _____

E.P.A. Reg. No. 1269-56

DE WITT CHEMICAL COMPANY

Camera's Note
Poor Copy

1269-56
ACCEPTED
OCT 31 1972
FORWARDED TO THE
ED UNDER THE
SUBJECT
TO ATTACHED COMMENTS.