

WEEDS

Use Low Rate

- Beggartick
- Carpetweed
- Cocklebur
- Corn crowfoot
- Dodder
- Fleabane
- Galinsoga
- Giant ragweed
- Hemp
- Henbit
- Jimson weed
- Morning Glory (annual)
- Mustard
- Pigweed
- Prostrate vervain
- Poverty weed
- Ragweed
- Rape
- Shepherd's-purse
- Sticktight
- Buckhorn plantain
- Bull thistle
- Burdock
- Buttercups
- Hedge bindweed
- Loco weed
- Pennywort
- Plantains
- Roadside thistle
- Spanish needles
- Spiderwort
- Sunflowers (Perennial)
- Vervains

Use Higher Rate

- Bedstraw
- Chickweed
- Dogfennel
- Goosefoot
- Knotweed
- Lamb's-quarters
- Mallow (annual)
- Marestail
- Prickly lettuce
- Russian thistle
- Smartweed
- Spotted spurge
- Wild buckwheat
- Wild lettuce
- ~~Wild radish~~
- Western ragweed
- Bur ragweed
- Button weed
- Field bindweed
- Goatsbeard
- Goldenrod
- Oxeye daisy
- Pikeweed
- ~~Water hyacinth~~
- Yarrow

For smaller jobs, mix 1 ounce of this product with one gallon of water.

For the control of Wild Onions or Garlic, mix one gallon of this product in 50 gallons of water plus 1/4 pound of household detergent (or 2 1/2 ounces per gallon of water plus 1/3 ounce of detergent) and apply in spring or fall at the rate of 50 gallons per acre. (One gallon per 870 square feet.) Since Wild Onions and Garlic bulbs and bulblets remain dormant in the soil for years, this treatment should be repeated at least 3 times. Spring-Fall-Spring or Fall-Spring-Fall. NOTE: 2,4-D and 2,4,5-T will damage creeping grasses such as bent, and should not be applied to this variety of turf. It will also damage clover and other legumes. Do not apply to pastures unless injury to clover and other legumes can be tolerated.

**CAUTION**

Do not permit spray to come in contact with any valuable ornamental or crop plants, for it will kill them. Course sprays are less likely to drift than are fine sprays.

DO NOT contaminate irrigation ditches or drinking water. Avoid spray drift to susceptible plants. Do not use on newly seeded lawns or turf. Do not graze treated areas by dairy animals within 7 days after treatment.

After applying material, DO NOT use this sprayer for subsequent application of insecticides or fungicides, for slight residue could damage valuable ornamental plants.

DO NOT store near seeds, fertilizer, insecticides, or fungicides.

DO NOT take internally. Avoid contact with skin, eyes, and clothing.

Contents are guaranteed to conform to ingredients statement. Seller makes no warranty of any kind, express or implied, concerning use of this product. Buyer assumes all risk of use or handling, whether in accordance with directions or not. KEEP OUT OF THE REACH OF CHILDREN.

Net Contents: \_\_\_\_\_ Gals.

USDA REG. NO. 1269-33

[1269-102] 2 pgs

Rev. 4-25-69  
Front Panel

~~732 H-5668~~

DEWITT 14D69

One-Shot Aerosol

Non-Toxic to Humans and Pets  
Kills Both Flying and Crawling Insects

ACCEPTED  
MAY 19 1969  
UNDER THE FEDERAL INSECTICIDE ACT  
FOR ECONOMIC PEST CONTROL  
ED UNDER NO. 1269-102 SUBJECT  
TO ATTACHED COMMENTS.

Effective against flies, mosquitoes, gnats, flying moths, roaches, ants, and silverfish.

May be used for intermittent spraying or as a total release aerosol.

ACTIVE INGREDIENTS: 14.5%

Pyrethrins	0.3%
Technical Piperonyl Butoxide*	0.6%
N-octyl bicycloheptene dicarboximide	1.0%
Petroleum Distillate	12.6%

INERT INGREDIENTS: 85.5%

\* Consists of 0.48% (butylcarbityl) (6 propylpiperonyl) ether and 0.12% related compounds.

CAUTION: Keep Out of Reach of Children. See further precautions on side panel.

NET CONTENTS 6 Oz. Av.

Side Panel

RT 5-19-69

~~HIGHLY CONCENTRATED~~ FOR COMMERCIAL AND INDUSTRIAL USE.

*In whole and area of food processing plant,* DIRECTIONS

For Flying Insects: Close doors and windows. Foods should be removed or covered during treatment. \* All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product as a space spray in these areas, apply only when plant is not in operation. One 6-ounce can will treat 16,000 cubic feet. Place aerosol on table or other support near center of area to be treated. Depress actuator and turn cap in direction of arrow to lock. Entire contents will be dispensed by this method.