

PM 32 1270-24 4  
12-31-77

**Harmful If Swallowed:** If it is, Use The Following Antidotes: Give milk, raw egg white, mucilage, gruel, or cornstarch paste, and follow with an emetic (tablespoonful of mustard in a tumbler of warm water). CALL A PHYSICIAN. If concentrated disinfectant comes in contact with the skin, flood with water and bathe with diluted alcohol. Causes eye and skin irritation. Do not get in eyes or on skin. In case of contact, flush with plenty of water for 15 minutes. If eye irritation persists, get medical attention. Avoid contamination of food.

### STORAGE AND DISPOSAL

Store in a cool, dry place away from heat and open flame. Keep container closed when not in use. DO NOT contaminate water, food, or feed by storage and disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Small plastic containers, after rinsing and puncturing as above, may be disposed of by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

This product is effective against the following organisms:

Salmonella typhosa	Streptococcus pyogenes
Shigella paratyphosae	Mycobacterium tuberculosis
Pseudomonas aeruginosa	Trichophyton interdigitale
Mycobacterium smegmatis	Salmonella schottmuelleri
Staphylococcus aureus	Escherichia coli

# ZEP-O-MINT

**A FRAGRANTLY SCENTED GENERAL  
PURPOSE DISINFECTANT  
GERMICIDE-DISINFECTANT  
TUBERCULOCIDE-FUNGICIDE  
BACTERICIDE-DEODORANT**

PHENOL COEFFICIENTS (AOAC METHOD)

S. aureus	11
S. typhosa	18
Pseudomonas aeruginosa	2

Passes AOAC First Action Procedure for determining Tuberculocidal Activity vs. Mycobacterium tuberculosis Stain H37Rv at a dilution of 1:64. Contains a broad spectrum germicide which works against a wide variety of pathogenic organisms. Deodorizes as it disinfects.

**ACTIVE INGREDIENTS:**

o-Benzyl-p-chlorophenol ..... 5.0%

**INERT INGREDIENTS:** ..... 95.0%

TOTAL ..... 100.0%

## WARNING

**KEEP OUT OF REACH OF CHILDREN**

See Left Side Panel for Additional Precautionary Statements and Statement of Practical Treatment.

E.P.A. REG. NO. 1270-24

E.P.A. EST. NO. 1270-GA-1

MANUFACTURED BY  
**ZEP MANUFACTURING COMPANY**  
DIVISION OF NATIONAL SERVICE INDUSTRIES, INC.  
ATLANTA, GEORGIA 30301

### DIRECTIONS

This is a concentrate. Use dilute as indicated below.

**As a General Disinfectant:** Be sure that all dirt is swept up and removed before applying, and that disinfectant solution comes into contact with all contaminated areas. To disinfect floors, cellars, basements, toilet rooms, bathrooms, lavatories, urinals, sinks, etc., clean thoroughly; use 1 ounce of this preparation to every gallon of water; mop in regular manner. If it is impractical to mop specific places, use 2 ounces per gallon of water and spray thoroughly over entire surface. Allow 2 minutes contact time.

To disinfect kitchen tables, drainboards, marble, tile, linoleum, limestone, glass, rubber, syrup cans, soda fountains, refrigerators, cooling rooms, refrigerator cans, garbage cans, rinse thoroughly with a dilution of 1 ounce of this product to a gallon of water after cleansing in regular manner, allowing contact for 5 minutes. All treated areas which will subsequently contact food products must be thoroughly rinsed with potable water prior to reuse.

For more drastic conditions such as disinfection of exceptionally dirty urinals, garbage cans, etc., a most thorough cleaning must be done by use of strong cleaning solutions, removing all dirt, then make sure disinfectant comes in contact with all contaminated areas; disinfect with a solution of 2 ounces of disinfectant per gallon and allow 2 minutes contact time.

To disinfect athlete's food fungi in locker rooms, prepare a solution of 1-1/2 ounces per gallon of water and thoroughly wet locker room surfaces, such as floors, walls, benches, showers, with this mixture. Deodorizes while it disinfects.

**FOR INDUSTRIAL AND INSTITUTIONAL USE ONLY**

PROD. #1835

190A

NET CONTENTS/GALS.

CONTROL NO.

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.