


Rev. 4-5-68
ALL ONE PANEL



ZEP FORMULA 31

~~DEVELOPED BY ZEP~~

For Control of Rats and Mice -- Contains *Fumarin

READY-TO-USE

For use in farms, dairies, poultry yards, cribs, warehouses, grain elevators, feed mills, out-buildings, dumps, wharfs, boats.

ACTIVE INGREDIENTS:

3-(Alpha-Acetyl-furfuryl)-4-Hydroxycoumarin* 0.025%

INERT INGREDIENTS:

99.975%

*Fumarin's Trademark Registered by Amchem Products, Inc.

DIRECTIONS

For Rats: Use one or two bait placements (one-fourth to one pound each) at every point of large activity. Look for such tell-tale signs as grawings, tracks or trails, burrows, drippings, or damage. An uninterrupted supply of bait should be maintained for at least 10 days. Bait should be placed in locations not accessible to children, pets, and domestic animals, or in tamper-proof boxes.

For Mice: Tablespoon amounts (one-fourth to one-half ounce) of bait should be placed at 8 to 12 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days. Bait should be placed in locations not accessible to children, pets, and domestic animals, or in tamper-proof boxes.

GENERAL INFORMATION

Place material in a protected spot where the rats and mice ordinarily feed and water, or where they travel or live. Baits should be placed along walls and in corners rather than in the middle of the room.

Use shallow containers for the bait. Protect CHILDREN, PETS, and DOMESTIC ANIMALS from bait.

Keep feeding stations well supplied. Inspect and replenish each station daily for the first week and regularly thereafter until bait is no longer being taken. Never allow a feeding station to remain empty for over 48 hours. If some bait stations are untouched, move to a different location. If baits become sour or moldy, as might be the case during extended periods of hot and humid weather, replace with fresh bait as needed.

Where re-infestation from dumps and fields may occur, permanent stations should be set out and replenished as needed.

PROVEN! ECONOMICAL! EFFECTIVE!

A new chemical for effective rodent control. Contains an anticoagulant chemical. Its remarkable killing power depends on its being consumed regularly in small doses for a period of days -- usually five to fourteen.

Has unusual properties which make this technique practical: No bait shyness -- Odorless and tasteless -- No pre-baiting -- Inexpensive and easy to use -- No tolerance buildup.

CAUTION

-KEEP OUT OF REACH OF CHILDREN-

Harmful if swallowed. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions may be indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

Avoid contamination of feed and foodstuffs.

NET CONTENTS/Lbs.

USDA Reg. No. 1270-63

~~1270-63~~

Manufactured by

ZEP MANUFACTURING COMPANY

Atlanta, Georgia