

ZEP FORMULA 2964

~~DISINFECTANT-DETERGENT~~

Disinfectant-Detergent

For Cleaning and Disinfecting Poultry and Livestock
Houses and Equipment


ACTIVE INGREDIENTS: 72.00%

Anhydrous sodium carbonate	60.00%
Potassium dichloro-s-triazinetriene	10.00%
Sodium dodecyl benzene sulfonate	2.00%

INERT INGREDIENTS: 28.00%

Anhydrous sodium tripolyphosphate
Sodium sulfate

Directions for Use



All tanks and spraying equipment should be rinsed with water after application. Use adequate ventilation in enclosed houses. If applied with a high pressure sprayer, use a face mask. Use hot or cold water for dilution.

POULTRY HOUSES, LIVESTOCK PENS, EQUIPMENT - To Clean and Disinfect: For hard, non-porous surfaces, use one ounce per gallon (provides 400 ppm available chlorine); for porous and hard-to-clean surfaces, use one-and-a-half ounces per gallon (provides 600 ppm available chlorine).

Remove debris, litter, and loose dirt. Treat all exposed areas, floors, walls, ceilings, posts, and support beams, scrubbing or applying ZEP FORMULA 2964 ~~DISINFECTANT-DETERGENT~~ solution with a pressure sprayer. Allow houses and pens to dry before re-introducing poultry or livestock.

Clean all equipment, such as feeders, brooders, waterers, with a solution of ZEP FORMULA 2964 ~~DISINFECTANT-DETERGENT~~. Allow to dry without rinsing.

(Continued)

DRINKING FOUNTAINS FOR LIVESTOCK AND POULTRY - To be Cleaned and Sanitized Daily: Scrub or dip in a solution of ZEP FORMULA 2964 ~~2964~~ until clean, using a concentration of one-half ounce per gallon (provides 200 ppm available chlorine). Do not rinse. Refill with fresh water.

CLEANING AND DISINFECTING IN HATCHERIES: For hard, non-porous surfaces, use one ounce per gallon (provides 400 ppm available chlorine); for porous and hard-to-clean surfaces, use one-and-a-half ounces per gallon (provides 600 ppm available chlorine).

Twice a week or more often, if necessary, scrub or clean with a pressure sprayer all hatchery walls, ceilings, and floors, using a solution of ZEP FORMULA 2964 ~~2964~~. Do not rinse.

Remove all trays and equipment from incubators and clean out debris. Scrub with a solution of ZEP FORMULA 2964 ~~2964~~. Allow to dry without rinsing. Clean dirt and debris from trays. Wash with a solution of ZEP FORMULA 2964 ~~2964~~. Rinsing is not necessary.

CAUTION

-KEEP OUT OF REACH OF CHILDREN-

Harmful if swallowed. Prolonged contact may irritate skin, eyes, or nose. Avoid exposure to the spray mist which may cause temporary irritation of the eyes or respiratory tract. If taken internally, feed milk. Call a physician.

USDA Reg. No. 1270-
~~1270~~

NET CONTENTS: _____ Gals.

7/13/67

ACCEPTED
Aug 22, 1967
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROVIDENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 1270-112

Comments

ZEP FORMULA 2964

~~DEWITT 10064~~

Disinfectant-Detergent

For Cleaning and Disinfecting Poultry and Livestock
Houses and Equipment

ACTIVE INGREDIENTS: 72.00%

Anhydrous sodium carbonate	60.00%
Potassium dichloro-s-triazinetrione	10.00%
Sodium dodecyl benzene sulfonate	2.00%

INERT INGREDIENTS: 28.00%

Anhydrous sodium tripolyphosphate
Sodium sulfate

Directions for Use

All tanks and spraying equipment should be rinsed with water after application. Use adequate ventilation in enclosed houses. If applied with a high pressure sprayer, use a face mask. Use hot or cold water for dilution.

POULTRY HOUSES, LIVESTOCK PENS, EQUIPMENT - To Clean and Disinfect: For hard, non-porous surfaces, use one ounce per gallon (provides 400 ppm available chlorine); for porous and hard-to-clean surfaces, use one-and-a-half ounces per gallon (provides 600 ppm available chlorine).

Remove debris, litter, and loose dirt. Treat all exposed areas, floors, walls, ceilings, posts, and support beams, scrubbing or applying ZEP FORMULA 2964 ~~DEWITT 10064~~ solution with a pressure sprayer. Allow houses and pens to dry before re-introducing poultry or livestock.

Clean all equipment, such as feeders, brooders, waterers, with a solution of ZEP FORMULA 2964 ~~DEWITT 10064~~. Allow to dry without rinsing.

(Continued)

BEST DOCUMENT AVAILABLE

ACCEPTED

AUG 22 1967

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND FUNGICIDE ACT
FOR ECONOMIC PESTICIDE REGISTRATION
ED UNDER SECTION 241. SUBJECT
TO ATTACHED COMMENTS.

DRINKING FOUNTAINS FOR LIVESTOCK AND POULTRY - To be Cleaned and Sanitized Daily: Scrub or dip in a solution of ZEP FORMULA 2964 ~~DEWIT-10064~~ until clean, using a concentration of one-half ounce per gallon (provides 200 ppm available chlorine). Do not rinse. Refill with fresh water.

CLEANING AND DISINFECTING IN HATCHERIES: For hard, non-porous surfaces, use one ounce per gallon (provides 400 ppm available chlorine); for porous and hard-to-clean surfaces, use one-and-a-half ounces per gallon (provides 600 ppm available chlorine).

Twice a week or more often, if necessary, scrub or clean with a pressure sprayer all hatchery walls, ceilings, and floors, using a solution of ZEP FORMULA 2964 ~~DEWIT-10064~~. Do not rinse.

Remove all trays and equipment from incubators and clean out debris. Scrub with a solution of ZEP FORMULA 2964 ~~DEWIT-10064~~. Allow to dry without rinsing. Clean dirt and debris from trays. Wash with a solution of ZEP FORMULA 2964 ~~DEWIT-10064~~. Rinsing is not necessary.

CAUTION


-KEEP OUT OF REACH OF CHILDREN-

Harmful if swallowed. Prolonged contact may irritate skin, eyes, or nose. Avoid exposure to the spray mist which may cause temporary irritation of the eyes or respiratory tract. If taken internally, feed milk. Call a physician.

USDA Reg. No. 1270-
~~10000~~

NET CONTENTS: _____ Gals.

7/13/67



BEST DOCUMENT AVAILABLE
Comments