

ACCEPTED
 FEB 12 1969
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND ROACENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 ED UNDER NO. *17012a* SUBJECT
 TO ATTACHED COMMENTS.

Rev. 2-3-69

ZEPONA

~~XXXXXXXXXX~~

Insecticide

A specially prepared insecticide for control of resistant and non-resistant insects.

ACTIVE INGREDIENTS: 100%

2,2-dichlorovinyl dimethyl phosphate	0.46%
Related compounds	0.04%
Petroleum hydrocarbons	99.50%

USES

For use in industrial plants, inedible products area of food processing plants, outdoor areas, in horse barns, dairy barns, milksheds, dog kennels, theaters, warehouses, and restaurants.


Controls flies, gnats, mosquitoes, wasps, fruit flies, flying moths, cockroaches, millipedes, centipedes, ants, spiders, silverfish, fleas, and brown dog ticks.

DIRECTIONS

Fogging Applications:

To kill insects such as flies, wasps, fruit flies, and flying moths in industrial plants, in inedible products areas of food processing plants (DO NOT USE IN EDIBLE PRODUCTS AREAS), and in warehouses, apply as a dense fog enveloping the area at a rate of 1 quart per 8,000 cubic feet. When treating inedible products areas of food processing plants, precautions should be taken to avoid contamination of packaged or stored foods. Before fogging, reduce air movement as much as possible by closing doors, windows, etc., and turning off fans. All electrical connections should be eliminated as well as other sources of ignition. Allow 5 to 10 minutes after fogging before ventilating area. Keep unprotected persons out of the operational area.

To kill flies, mosquitoes, wasps, gnats, flying moths, and other insects which prove troublesome around picnic areas, playgrounds, patios, and other outdoor areas, apply as a dense fog using a thermal fogging device such as the Aero-Dyne Fogger. Material should be fogged at a rate of 1½ to 2½ gallons of insecticide per acre. It should be applied at a time of low wind velocity, and fog should always be directed downwind from the operator and fogging machine. Fog should be applied in such a manner that it will drift over the infested areas slowly as possible. Keep unprotected persons out of the operational area.



Applications as a Surface Spray:

For control of border to control insects, such as cockroaches, spiders, millipedes, centipedes, ants, silverfish in restaurants, theaters, food processing plants, industrial plants, warehouses, and wherever else they infest, apply as a surface spray at the rate of 1 quart per 1,000 square feet. Pay particular attention to breeding and nesting sites such as cracks in walls, along baseboards, in closets, storage cabinets. Do not contaminate food, food handling equipment, or drinking water. Do not apply in the edible products areas of food processing plants.

To kill annoying flies and fly maggots in and around horse barns, dairy barns, and milksheds, apply as a wet spray using a low pressure sprayer. Surfaces such as doorways, feed storage room floors and walls, alleys, window sills should be thoroughly wetted with the spray at a rate of 1 quart per 1,000 square feet. Do not contaminate milk, milking utensils, feed, or water.

For control of maggots in breeding areas such as manure piles, garbage dumps, apply as a coarse spray at a rate of 1 to 2 quarts per 100 square feet, thoroughly wetting infested surfaces.

To kill flies, gnats, and other annoying insects around dog kennels, apply as a spray, 1 quart per 1,000 square feet. Apply to such surfaces as curtains, runways, window sills, and ledges. To control fleas and head dog ticks, increase dosage to 2 quarts per 1,000 square feet.

As a face spray for cattle to control flies and gnats, apply 1 to 2 fluid ounces per animal, as a fine mist spray, daily with hand or automatic sprayer. Do not get into animal's eyes. Apply spray so as to wet the hair. Do not apply in excess of 2 fluid ounces per animal per day.

(2) (b) (3)

REGISTRATION OF PESTICIDES

Avoid contact with skin and eyes. Avoid breathing spray mist. May be fatal if swallowed, inhaled, or absorbed through the skin. Wash thoroughly with soap and water after use and before eating or smoking.

Do not contaminate food, water, or food containers. Avoid use in poultry.

This product is toxic to fish and wild birds. Keep out of lake, stream, or pond. Apply this product only as directed on this label. Birds feeding on treated areas may be killed.

Do not re-use empty drum. Return to drum receiver or destroy by puncturing or crushing and burying in a safe place.

Do not use in edible products areas of food processing plants.

USDA Reg. No. _____

NET CONTENTS/GALS.

Manufactured By

REP MANUFACTURING COMPANY

Atlanta, Georgia

ACCEPTED
Feb-12-1969
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PESTICIDE REGISTERED
UNDER NO. 1370720