

For General Vegetation


NO-GRO GRANULAR will give quick knock-down of vegetation and residual control of broad-leaved weeds and grasses on parking lots, fence lines, highway and railway sites, drainage ditches and other similar areas. For dry granular treatment, apply 1/2 to 2 lbs. per 100 square feet. The lighter rate should only be used at time of weed emergence on moist soil or when precipitation is anticipated while weeds are still young.

Persistent perennial weeds are most effectively treated when nearing maturity or during dormant stages in the fall. Spring applications may also be made under moderate rainfall conditions. Summer applications are recommended only when rainfall is dependable for leaching the herbicide into the soil. Apply 2 to 4 lbs. per 100 sq. ft. in the dry granular form. On deep rooted perennials such as bindweed, Canada thistle, Bermuda grass, and Quack grass, the heavier rates may be required.

EPA Reg. # 1325-75

100 GALLONS

GALLONS



NO-GRO Granular

GRANULAR, NONSELECTIVE GRASS AND WEEDKILLER

Active Ingredients

Sodium Chlorate (NaClO_3) 30%
 Sodium Metaborate Tetrahydrate ($\text{Na}_2\text{B}_4\text{O}_7 \cdot 4\text{H}_2\text{O}$) . . . 63%
 (Boron Trioxide (B_2O_3) Equivalent 23.3%)

Inert Ingredients

80% of 100%

CAUTION

KEEP OUT OF REACH OF CHILDREN *3/8/75*

For use as a lawn and garden maintenance treatment only. Do not use for any other purpose. For domestic purposes, a solution diluted with water may be used for household cleaning. Do not use on equipment, machinery, or vehicles. Do not use on plants or animals, and do not use on trees, shrubs, or ornamentals until the foliage is reached through the canopy. Do not use chemical equipment that should be thoroughly cleaned after use. Do not store in direct contact with soil and/or concrete floor.

Do not reuse container. Destroy when empty

Sold by



NO GRO GRA
 rooted perer
 Dallis grass
 lines and
 pounds per
 Johnsongras
 late fall.

FOI

For use as
 asphalt pavi
 paving due t
 per 100 squa
 mended
 For DRY tre
 application,
 100 square f

This materia
 be toxic to
 treated area
 or more yea
 use or appli
 be treated ar
 shrubs, crop
 not intended
 precautions
 by roots of
 applications
 off to areas
 which will l
 injury to d
 equipment or
 local condit
 culture. eq
 weed specta
 to fit local