


50 LBS. NET WEIGHT
(22.7 KILOS)

**WALBRIDGE
LABORATORIES
PLANTING CORPORATION**

ACCEPTED
OCT 24 1983
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under: **EPA Reg. No. 1275-1**

Basic Copper Sulfate



REST DOCUMENT AVAILABLE

KEEP OUT OF REACH OF CHILDREN
CAUTION
HARMFUL IF SWALLOWED
AVOID CONTACT WITH EYES
In case of contact with eyes or face, flush with plenty of water. Contact with skin, wash with soap and water. See label for first aid instructions.

NOT FOR MEDICINAL USE

EPA REG. NO. 1275-1
EPA LABELING NO. 1772-1A8

ACTIVE INGREDIENT:
COPPER SULFATE MONOHYDRATE METALLIC 52%
INERT INGREDIENTS: 48%
TOTAL 100%

MANUFACTURED BY:



See label for first aid instructions and precautions for use. For more information, contact the manufacturer, Walbridge Laboratories, Inc., 10000 West 10th Street, Denver, Colorado 80231. For more information, contact the State Agricultural Experiment Station, University of California, Davis, California 95616.

Basic Copper Sulphate

BASIC COPPER SULPHATE may be used on crops for which liquid mixtures are recommended (in some cases it should be added to the recommended liquid mixture). Use Basic Copper Sulphate at one-half the amount of copper sulfate (blue stone) required in a recommended Bordeaux mixture.

Amount of BASIC COPPER SULPHATE Required to Equal an 8-2-10 Bordeaux Mixture

Yield of Crop per Acre	100 lbs	250 lbs	500 lbs	1000 lbs	2000 lbs	3000 lbs	4000 lbs
Basic Copper Sulphate	4 lbs	2 1/2 lbs	1 lb	5/8 lb	3/4 lb	5/8 lb	5/8 lb

DIRECTIONS FOR USE

All directions are to be printed in Copper Sulphate per 100 gallons of water. For concentrated sprays, use the amount of Basic Copper Sulphate per 100 gallons of water. Consult with your State Agricultural Extension Agent for spray recommendations for your area.

FRUITS AND VEGETABLES

APPLES - Apply to the leaves of the tree when the leaves are first out. Repeat every 10 to 15 days until the fruit is ready to harvest. Use 4 lbs per 100 gallons of water.

PEACHES - Apply to the leaves of the tree when the leaves are first out. Repeat every 10 to 15 days until the fruit is ready to harvest. Use 4 lbs per 100 gallons of water.

PLUMS - Apply to the leaves of the tree when the leaves are first out. Repeat every 10 to 15 days until the fruit is ready to harvest. Use 4 lbs per 100 gallons of water.

CHERRIES - Apply to the leaves of the tree when the leaves are first out. Repeat every 10 to 15 days until the fruit is ready to harvest. Use 4 lbs per 100 gallons of water.

ORCHARD FRUIT TREES - Apply to the leaves of the tree when the leaves are first out. Repeat every 10 to 15 days until the fruit is ready to harvest. Use 4 lbs per 100 gallons of water.

VEGETABLES - Apply to the leaves of the plant when the leaves are first out. Repeat every 10 to 15 days until the vegetable is ready to harvest. Use 4 lbs per 100 gallons of water.

GRAPES - Apply to the leaves of the vine when the leaves are first out. Repeat every 10 to 15 days until the fruit is ready to harvest. Use 4 lbs per 100 gallons of water.

STRAWBERRIES - Apply to the leaves of the plant when the leaves are first out. Repeat every 10 to 15 days until the fruit is ready to harvest. Use 4 lbs per 100 gallons of water.

SPRING GREENS - Apply to the leaves of the plant when the leaves are first out. Repeat every 10 to 15 days until the vegetable is ready to harvest. Use 4 lbs per 100 gallons of water.

SPINACH - Apply to the leaves of the plant when the leaves are first out. Repeat every 10 to 15 days until the vegetable is ready to harvest. Use 4 lbs per 100 gallons of water.

BEANS - Apply to the leaves of the plant when the leaves are first out. Repeat every 10 to 15 days until the vegetable is ready to harvest. Use 4 lbs per 100 gallons of water.

PEAS - Apply to the leaves of the plant when the leaves are first out. Repeat every 10 to 15 days until the vegetable is ready to harvest. Use 4 lbs per 100 gallons of water.

MIXING DIRECTIONS

1. Measure the amount of Basic Copper Sulphate required for the area to be treated.

2. Add the Basic Copper Sulphate to the water in the sprayer.

3. Mix thoroughly before using.

PRECAUTIONARY STATEMENTS

CAUTION

Basic Copper Sulphate is a toxic substance. It is harmful to fish and other aquatic life. It is also harmful to birds and mammals. Do not use in or near water bodies. Do not use on lawns or ornamental plants. Do not use on children's play areas. Do not use on crops that are eaten raw.

STORAGE AND DISPOSAL

Store in a cool, dry place. Do not store in a metal container. Do not store in a container that has been used for other chemicals. Do not store in a container that is damaged or leaking. Do not store in a container that is not clearly labeled.

CONTAINER DISPOSAL

Do not reuse the container. Do not burn the container. Do not dump the container in the trash. Do not dump the container in the water. Do not dump the container in the soil.