

FIFTY

GALLONS

DIRECTIONS

Fogging or Space Spraying: To kill Flies, Mosquitoes, Gnats and Spiders in dairy and beef cattle barns, apply daily as a fog or mist inside building at rate of 1 pint per 8,000 cubic feet. Keep cattle in barn when spraying. Let spray settle before milking. Do not apply in areas where animals have received direct applications of this product within 8 hours.

Direct Application: To protect dairy and beef cattle from Face Flies, Houseflies, Stable Flies, Horn Flies, Mosquitoes and Gnats, spray animal daily using 1 to 2 fluid ounces per animal. Apply as a mist spray so as to wet the hair. Do not soak the skin or wet the hide. Pay particular attention to the face, back, flanks and legs. Experience indicates that two applications of 1 fluid ounce per animal morning and night give outstanding control of the above pests. Do not apply in excess of 2 fluid ounces per animal per day.

Additional Uses: Dilute one part McNess Fogger Oil and Dairy Spray with one part of fuel oil, diesel oil or kerosene. Apply this solution as a coarse wet spray at the rate of one quart per 1,000 square feet to control Flies, Gnats, Mosquitoes, Roaches, Spiders, Fleas, Ants, Sowbugs, Brown Dog Ticks.

For milk sheds, stables, livestock barns, loafing sheds, pig pens and outdoor areas, treat around doorways, feed storage rooms, alleyways and window sills.

McNess[®]

FOGGER OIL AND DAIRY SPRAY

For Killing Houseflies, Horn Flies, Stable Flies, Face Flies, Mosquitoes, Gnats, Spiders and certain Other Insects in Dairy Barns and Other Farm Buildings. Also for direct application to Dairy and Beef Cattle.

ACTIVE INGREDIENTS

*2, 2-Dichlorovinyl Dimethyl Phosphate	0.93%
*Related Compounds	0.07%
Petroleum Hydrocarbons	98.90%

INERT INGREDIENT

Epichlorohydrin	0.10%
	<u>100.00%</u>

*Equivalent to 1.00% Vapona[®] Insecticide

USDA Registration No. 1304-45

WARNING: KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTION ON BACK PANEL

Manufactured by
FURST-McNESS COMPANY
Chemists and Manufacturing Pharmacists
Established 1908

FREEPORT, ILL.
BALTIMORE, MD.

OAKLAND, CALIF.
MEMPHIS, TENN.

In poultry houses, apply to manure, window sills, outside of penned enclosures and on floor of feed storage rooms.

In treating dog kennels apply to outside dog runways, window sills and ledges. For control of fleas and brown dog ticks increase dosage to 2 quarts per 1,000 square feet.

For control of fly maggots in maggot breeding areas such as garbage dumps and manure piles apply at the rate of 2 quarts per 100 square feet.

CAUTION

Do not contaminate water, feed, milk or milking equipment. May be fatal if swallowed, inhaled or absorbed through the skin. Avoid prolonged breathing of spray mist. Wash thoroughly with soap and water after handling and before eating and smoking. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal.

Vapona is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticide or chemicals.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Apply this product only as specified on this label.

LABORATORY CONTROL
NO.

ACCEPTED
2/11/70
1304-45
FURST-McNESS COMPANY
OAKLAND, CALIF.