



A Combination Fungicide and Insecticide

NEW • IMPROVED • CONTAINS SEVIN*

Active Ingredients:

Maneb (Manganese ethylene bisdithiocarbamate) 3.9%

Carbaryl (1-Naphthyl N-Methylcarbamate) . . 5.0%

Inert Ingredients: (Including ZnSO₄) 91.1%

EPA Reg. No. 1304-61

EPA Est. No. 1304-IL-1

*Registered trade-mark of Union Carbide Corporation for Carbaryl Insecticide.

A Dual Purpose Product—Highly effective against listed insects and fungi. It is no longer necessary to purchase several different products to give good control in your garden. You can rely on McNess Garden Dust to safeguard your plants and shrubs effectively.

Economical to Use—Helps protect your investment in your flower and vegetable gardens and ornamental shrubs.

Ready to Use—Needs no mixing or diluting—simply use as it comes from the package.

Highly Effective—Kills listed insects and will control many of the most prominent destructive blights or fungus diseases.

Apply as soon as plants become infested with insects and damage is noticeable. Apply at 5 to 10 day intervals as needed. Dust thoroughly to get good coverage on upper and lower sides of leaves. Use a good duster. Dust adheres better in late evening or early morning when vegetation is moist with dew and when the air is calm.

CAUTION: KEEP OUT OF REACH OF CHILDREN

HARMFUL IF INHALED OR SWALLOWED. Do Not Breathe Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Skin Contact. Wash Thoroughly After Handling. Note for Physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal. Avoid contamination of food. Do not reuse empty container. Destroy when empty.

DS

Use McNess Garden Dust on:

BEANS—(Cowpeas, Blackeyed Peas)—for Downy Mildew. (Lima, Snap, Green)—for Anthracnose, Downy Mildew. (Cowpeas, Blackeyed Peas, Lima, Snap, Green)—for Mexican Bean Beetles, Bean Leaf Beetles, Cucumber Beetles, Flea Beetles, Japanese Beetles, Leafhoppers, Velvetbean Caterpillars, Western Bean Cutworms, Armyworms, Corn Earworms, Stink Bugs, Tarnished Plant Bugs. Dust plants thoroughly at first sign of disease or infestation. Repeat at 5 to 10 day intervals. **DO NOT APPLY WITHIN 7 DAYS OF HARVEST.**

CARROTS—for Cercospora Leaf Spot, Alternaria, Flea Beetles, Leafhoppers, Harlequin Bugs. Begin when plants are 6 weeks old. Repeat at 7 to 10 day intervals.

CABBAGE, BROCCOLI, BRUSSEL SPROUTS, CAULIFLOWER—for Alternaria Leaf Spot, Downy Mildew, Flea Beetles, Harlequin Bugs, Armyworms, Cabbage Worms and Loopers, Corn Earworms. Dust plants thoroughly at first sign of disease or infestation. Repeat at 3 to 7 day intervals. **DO NOT APPLY WITHIN 7 DAYS OF HARVEST.**

SWEET CORN—for Helminthosporium Leaf Blight, Corn Earworms, Corn Rootworm Adults, European Corn Borers, Fall Armyworms, Flea Beetles, Japanese Beetles, Sap Beetles and Leafhoppers. For larvae in whorl (budworm damage) and Foliage Feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ear, apply at 3-day intervals starting when first silks appear, and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are essential for effective control. Do not feed treated forage to dairy animals or to animals being finished for slaughter. Applications of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

MELONS—for Downy Mildew, Anthracnose, Cercospora Leaf Spot, Burning Stem Blight, Pickleworms, Melonworms, Flea Beetles, Leafhoppers. Begin at late bunching when vines begin to run or when disease or infestation first appears. Do not use on watermelons in Florida. **DO NOT APPLY WITHIN 5 DAYS OF HARVEST.**

CUCUMBERS—Downy Mildew, Anthracnose, Alternaria (Macrosporium) Leaf Spot, Angular Leaf Spot, Pythium Fruit Rot, Pickleworms, Cucumber Beetles, Flea Beetles, Leafhoppers. Begin when plants begin to run or when disease or infestation first appears. Repeat at 7 to 10 day intervals. **DO NOT APPLY WITHIN 5 DAYS OF HARVEST.**

PUMPKINS—for Downy Mildew, Angular Leaf Spot, Flea Beetles, Leafhoppers, Melonworms. Begin when disease or infestation first begins. Repeat at 7 to 10 day intervals.

SQUASH—for Downy Mildew, Anthracnose, Flea Beetles, Leafhoppers, Squash Bugs. Begin when plants start to run or when infestation first appears. Repeat at weekly intervals. **DO NOT APPLY WITHIN 5 DAYS OF HARVEST.**

GRAPES—for Black Rot, Grape Berry Moths, Japanese Beetles, June Beetles, Red-Banded Leaf Rollers, Grape Leaf Folders, Grape Leafhoppers and Grape Leaf Skeletonizers. Begin just before bloom for Black Rot. Apply as necessary. Apply just before first brood Leaf Folder larvae emerge from rolls and as needed for Leafhoppers.

COLLARDS, KALE—for Downy Mildew, Alternaria Leaf Spot, Flea Beetles, Harlequin Bugs, Leafhoppers, Armyworms, Cabbage Worms and Loopers, Corn Earworms, Spittlebugs, Tarnished Plant Bugs, Stink Bugs. Apply when disease or infestation first appears. Repeat at 7 to 10 day intervals. **DO NOT APPLY WITH 3 DAYS OF HARVEST.**

ENDIVE, LETTUCE—Downy Mildew, Flea Beetles, Harlequin Bugs, Leafhoppers, Armyworms, Cabbage Worms and Loopers, Corn Earworms, Spittlebugs, Tarnished Plant Bugs, Stink Bugs. Apply when disease or infestation first appears. Repeat at 7 to 10 day intervals. **DO NOT APPLY WITH 3 DAYS OF HARVEST.**

SPINACH—for Downy Mildew, White Rust, Flea Beetles, Harlequin Bugs, Leafhoppers, Armyworms, Cabbage Worms and Loopers, Corn Earworms, Spittlebugs, Tarnished Plant Bugs, Stink Bugs. Apply when disease or infestation first appears. Repeat at 7 to 10 day intervals. **DO NOT APPLY WITHIN 3 DAYS OF HARVEST.**

EGGPLANT—for Anthracnose, Early Blight (Alternaria), Phomopsis Blight or Fruit Rot, Flea Beetles, Leafhoppers, European Corn Borers, Fall Armyworms, Lace Bugs, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs. Apply when fruit starts to form or when infestation appears. Repeat at 7 to 10 day intervals.

PEPPERS—for Anthracnose, Cercospora Leaf Spot, Ripe Rot, Flea Beetles, Leafhoppers, European Corn Borers, Fall Armyworms, Lace Bugs, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs. Apply when disease or infestation first appears. Repeat at 7 to 10 day intervals.

TOMATOES—(Greenhouse and Field) for Cladosporium Leaf Mold, Early and Late Blight, Gray Leaf Spot (Tomphylium), Septoria Leaf Spot, Flea Beetles, Leafhoppers, European Corn Borers, Fall Armyworms, Lace Bugs, Tomato Fruit Worms, Tomato Hornworms, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs. Apply when fruit forms or infestation appears. Repeat at 7 to 10 day intervals. **DO NOT APPLY WITHIN 5 DAYS OF HARVEST.**

ROSES—for Black Spot, Cercospora Leaf Spot, Rust. Begin when first leaves unfold. Repeat at 7 to 10 day intervals.

GRASSHOPPERS on Above Crops—Dust as necessary.

BOX ELDER BUGS—Dust around foundation of house and a few feet of surrounding lawn.

Side Panels

ORNAMENTAL INSECT CONTROL

McNess Garden Dust can be safely used on a wide range of ornamental plants, flowers, shrubs and small trees, such as rose, carnation, gladiolus, chrysanthemum, lilac, arborvitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. Do not apply to Virginia Creeper or Boston Ivy. Injury to tender foliage may occur if plants are wet when treated or in presence of high humidity.

Apply to the upper and lower leaf surfaces of all foliage for complete coverage to control such pests as:

Apple Aphids
Bagworms
Birch Leaf Miners
Blister Beetles
Boxelder Bugs
Boxwood Leaf Miners

Elm Leaf Aphids
Elm Leaf Beetles
Flea Beetles
Gypsy Moths
Japanese Beetles
June Beetles

Lace Bugs
Leafhoppers
Mealy Bugs
Mimosa Webworms
Oak Leaf Miners

Orange Tortrix
Periodical Cicada
Plant Bugs
Puss Caterpillars
Psyllids
Rose Aphids

Rose-slug
Scale Insects
Tent Caterpillars
Thorn Bugs
Thrips (exposed)
Willow Leaf Bettles

NOTE

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Co-operative Agricultural Extension Service.

Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

ORNAMENTAL INSECT CONTROL

McNess Garden Dust can be safely used on a wide range of ornamental plants, flowers, shrubs and small trees, such as rose, carnation, gladiolus, chrysanthemum, lilac, arborvitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. Do not apply to Virginia Creeper or Boston Ivy. Injury to tender foliage may occur if plants are wet when treated or in presence of high humidity.

Apply to the upper and lower leaf surfaces of all foliage for complete coverage to control such pests as:

Apple Aphids
Bagworms
Birch Leaf Miners
Blister Beetles
Boxelder Bugs
Boxwood Leaf Miners

Elm Leaf Aphids
Elm Leaf Beetles
Flea Beetles
Gypsy Moths
Japanese Beetles
June Beetles

Lace Bugs
Leafhoppers
Mealy Bugs
Mimosa Webworms
Oak Leaf Miners

Orange Tortrix
Periodical Cicada
Plant Bugs
Puss Caterpillars
Psyllids
Rose Aphids

Rose-slug
Scale Insects
Tent Caterpillars
Thorn Bugs
Thrips (exposed)
Willow Leaf Bettles

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Co-operative Agricultural Extension Service.

Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

