

### DIRECTIONS FOR USE

**HOUSEHOLD INSECTS** (Roaches, Ants, Spiders, Silverfish, Crickets, Clover Mites, Earwigs) – In and about homes, dairies and food processing plants. Use 2% Malathion in water (5 ounces per gallon). Apply by means of coarse spray or paint brush to window sills, baseboards, drainboards, under sinks, stoves and to other areas frequented by insects. The 2% Malathion spray cannot be applied in the edible products areas of food processing plants.

**VEGETABLES** (Beans, Peas, Potatoes). For control of Aphids, use 1½ to 2 tsps. to 1 gallon water. Red Spider Mites, Mealy Bugs, White Fly, Leafhopper, Japanese Beetle Adults: 1 tablespoonful per gallon water. **NOTE:** Make no applications 7 days before harvest.

**ORNAMENTALS:** For the control of insects on ornamentals: Use 1½ pints per 100 gallons of water, making one or more applications as needed. For the control of Soft Brown Scale, Monterey Pine Scale and Black Scale Crawlers, use 2½ pints per 100 gallons of water, making one or more applications as needed. For control of Juniper Scale, Fletcher Scale, Bagworms and Oak Kermes, use 2 pints to 100 gallons of water.

**NOTE:** Do not use on Boston Maidenhair, Pteris Ferns, as well as species of Crassula. Do not apply on Petunias.

## MAGNOLIA

### 50% MALATHION

#### EMULSIFIABLE CONCENTRATE

Controls Certain Sucking and Chewing Insects

ACTIVE INGREDIENTS	% BY WT.
Malathion* .....	50%
Xylene .....	45%
INERT INGREDIENTS .....	5%
TOTAL .....	100%

\*O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
See other warnings on side panel.

When used as directed, controls Thrips, Aphids, Red Spider Mites, Garden Flea Hoppers, Japanese Beetles, Blister Beetles, Clover Mites, Two Spotted Mites, Mealy Bugs, Rose Chafers, Climbing Cut Worms, Web Worms, Tent Caterpillars, Grasshoppers, Crickets, and Ants on flowering and ornamental plants.

**FOR MOSQUITO CONTROL** – Mosquito larvae in standing water. Use 1 pint (½ lb. actual Malathion) per acre. Mix in sufficient water or oil to obtain even coverage when applied by air or ground equipment.

**HOUSE FLIES:** For reduction of fly population, use 2 gallons per 100 gallons of water, or 1 pint in 6 gallons of water for control in dairy barns, horse and sheep barns, poultry houses, manure piles, garbage heaps, city dumps.

**CAUTION:** Not to contaminate milk, milking equipment, feed or drinking water when applying the fly control treatments.

**CAUTION:** Harmful if swallowed. Avoid prolonged breathing of spray mist. Avoid repeated or prolonged contact with skin. Wash thoroughly. Avoid contamination of foodstuffs. Keep out of reach of children.

**FISH AND/OR WILDLIFE CAUTION:** This product is toxic to fish. Keep out of lakes, streams or ponds. Apply this product only as specified on this label.

**DO NOT REUSE EMPTY CONTAINER.** Destroy it by perforating or crushing. Bury or discard in a safe place.

**NOTE TO PHYSICIANS:** MALATHION UPON repeated prolonged use may cause cholinesterase inhibition. Atropine is antidotal.

**NOTICE OF WARRANTY:** Magnolia Chemical Co., Inc. makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label information.

U.S.D.A. Reg. No. 1307-24 Net Contents .....

4-24-70  
1307-24

## DIRECTIONS FOR USE

**HOUSEHOLD INSECTS** (Roaches, Ants, Spiders, Silverfish, Crickets, Clover Mites, Earwigs) – In and about homes, dairies and food processing plants. Use 2% Malathion in water (5 ounces per gallon). Apply by means of coarse spray or paint brush to window sills, baseboards, drainboards, under sinks, stoves and to other areas frequented by insects. The 2% Malathion spray cannot be applied in the edible products areas of food processing plants.

**VEGETABLES** (Beans, Peas, Potatoes). For control of Aphids, use 1½ to 2 tsps. to 1 gallon water. Red Spider Mites, Mealy Bugs, White Fly, Leafhopper, Japanese Beetle Adults: 1 tablespoonful per gallon water. **NOTE:** Make no applications 7 days before harvest.

**ORNAMENTALS:** For the control of insects on ornamentals: Use 1½ pints per 100 gallons of water, making one or more applications as needed. For the control of Soft Brown Scale, Monterey Pine Scale and Black Scale Crawlers, use 2½ pints per 100 gallons of water, making one or more applications as needed. For control of Juniper Scale, Fletcher Scale, Bagworms and Oak Kermes, use 2 pints to 100 gallons of water.

**NOTE:** Do not use on Boston Maidenhair, Pteris Ferns, as well as species of Crassula. Do not apply on Petunias.

# MAGNOLIA 50% MALATHION

## EMULSIFIABLE CONCENTRATE

Controls Certain Sucking and Chewing Insects

ACTIVE INGREDIENTS	% BY WT.
Malathion* .....	50%
Xylene .....	45%
INERT INGREDIENTS .....	5%
TOTAL .....	100%

\*O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
See other warnings on side panel.

When used as directed, controls Thrips, Aphids, Red Spider Mites, Garden Flea Hoppers, Japanese Beetles, Blister Beetles, Clover Mites, Two Spotted Mites, Mealy Bugs, Rose Chafers, Climbing Cut Worms, Web Worms, Tent Caterpillars, Grasshoppers, Crickets, and Ants on flowering and ornamental plants.

**MAGNOLIA CHEMICAL CO., INC.**  
P. O. BOX 20179  
2646 BODINEY LANE  
DALLAS, TEXAS  
247-7111

**FOR MOSQUITO CONTROL** – Mosquito larvae in standing water. Use 1 pint ½ lb. actual Malathion per acre. Mix in sufficient water or oil to obtain even coverage when applied by air or ground equipment.

**HOUSE FLIES:** For reduction of fly population, use 2 gallons per 100 gallons of water, or 1 pint in 6 gallons of water for control in dairy barns, horse and sheep barns, poultry houses, manure piles, garbage heaps, city dumps.

**CAUTION:** Not to contaminate milk, milking equipment, feed or drinking water when applying the fly control treatments.

**CAUTION:** Harmful if swallowed. Avoid prolonged breathing of spray mist. Avoid repeated or prolonged contact with skin. Wash thoroughly. Avoid contamination of foodstuffs. Keep out of reach of children.

**FISH AND OR WILDLIFE CAUTION:** This product is toxic to fish. Keep out of lakes, streams or ponds. Apply this product only as specified on this label.

**DO NOT REUSE EMPTY CONTAINER.** Destroy it by burning, foraging or crushing. Buy and discard in a safe place.

**NOTE TO PHYSICIANS:** MALATHION UPON repeated prolonged use may cause cholinesterase inhibition. Atropine is antidote.

**NOTICE OF WARRANTY:** Magnolia Chemical Co. makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label information.

U.S.D.A. Reg. No. 1307-24 Net Contents .....

4-24-70  
1307-24