

MAL-MAR

GRAIN PROTECTANT

AND

BIN SPRAY

*Direct Application
to grain in bulk or
as a top dressing.*

Contains 5 Lbs. Premium Grade Malathion Per Gal.

ACTIVE INGREDIENTS:

Malathion (0, 0-dimethyl phosphorothioate of diethylmercaptosuccinate)	57%
Aromatic petroleum hydrocarbons	33%
INERT INGREDIENTS	10%

**CAUTION: Keep out of reach of children.
See side panel for additional cautions.**

MALMOR COMPANY, INC.
2000 W. 10TH AVE
DALLAS, TEXAS
CN7-7111

DIRECTIONS FOR USE

1. DIRECT APPLICATION:

FOR THE PROTECTION OF STORED GRAINS such as wheat, oats, rice, corn, rye, barley, grain sorghum, and field or garden seeds, against confused flour beetle, rice weevil, granary weevil, saw-toothed grain beetle, flat grain beetle, red flour beetle, rusty grain beetle, lesser grain borer, and Indian meal moth, apply 1 pint of MAL-MAR in 2 to 5 gallons of water per 1000 bushels. Apply as the grain is being loaded or turned into final storage.

FIELD AND GARDEN SEEDS can be protected against the above grain pests with a dosage of 1/2 pint MAL-MAR in 1 to 2 1/2 gallons of water per 500 bushels of seed.

Do not use for food, feed or oil purposes.

2. TOP DRESSING:

TO PROTECT STORED GRAINS from attack by Indian meal moth, Angoumois grain moth, apply MAL-MAR to the surface of clean or uninfested grain at the rate of 1/2 pint in 1 to 2 gallons of water per 1000 square feet of grain surface area. Apply the spray evenly over the surface of the grain. Apply immediately after grain is loaded into storage and repeat if necessary.

U.S.D.A. Reg. No. 1307-55

DIRECTIONS FOR USE

3. RESIDUAL BIN SPRAY:

FOR A RESIDUAL WALL, FLOOR, AND MACHINERY SPRAY IN GRAIN ELEVATORS, in treating truck beds, box cars, and ships holds before loading grain, apply 1 gallon of MAL-MAR per 25 gallons of water making thorough application. Before applying spray, clean elevators, box cars, etc., thoroughly. Remove and burn all sweepings and debris.

CAUTION: Harmful if swallowed and absorbed through the skin. Avoid prolonged breathing of spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after using. Avoid contamination of feed or foodstuffs. There should be no application where milk is stored or processed, or in dwellings. Keep out of reach of children and domestic animals. Malathion, upon repeated or prolonged careless use, may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid. Atropine is antidotal.

Contents:

ED
f. 2. 1969
No. 111

MAL-MAR

GRAIN PROTECTANT

AND

BIN SPRAY

*Direct Application
to grain in bulk or
as a top dressing.*

DIRECTIONS FOR USE

1. DIRECT APPLICATION:

FOR THE PROTECTION OF STORED GRAINS such as wheat, oats, rice, corn, rye, barley, grain sorghum, and field garden seeds, against confused flour beetle, rice weevil, granary weevil, saw-toothed grain beetle, flat grain beetle, red flour beetle, rust grain beetle, lesser grain borer, and Indian meal moth, apply 1 pint of MAL-MAR in 1 to 5 gallons of water per 1000 bushels. Apply as the grain is being loaded or turned into final storage.

FIELD AND GARDEN SEEDS can be protected against the above grain pests with a dosage of 1/2 pint MAL-MAR in 1 to 2 1/2 gallons of water per 500 bushels of seed.

Do not use for food, feed or oil purposes.

2. TOP DRESSING:

TO PROTECT STORED GRAINS from attack by Indian meal moth, Angoumois grain moth, apply MAL-MAR to the surface of clean or uninfested grain at the rate of 1/2 pint in 1 to 2 gallons of water per 1000 square feet of grain surface area. Apply the spray evenly over the surface of the grain. Apply immediately after grain is loaded into storage and repeat if necessary.

U.S.D.A. Reg No. 1307-55

Contains 5 Lbs. Pesticide of Malathion Pesticide

ACTIVE INGREDIENTS

Malathion 5.0%
Diethyl phosphorodithioic acid
Non-hazardous

INERT INGREDIENTS

CAUTION: Keep out of reach of children.
See side panel for additional cautions.

MAGNOLIA CHEMICAL CO., INC.
P. O. BOX 20179
2646 RODNEY LANE
DALLAS, TEXAS
CH 7-7111

Malathion is a highly volatile liquid which will evaporate rapidly. It is highly toxic to fish and aquatic life. It is also highly toxic to birds and mammals. It is highly toxic to bees and other beneficial insects. It is highly toxic to cats and dogs. It is highly toxic to children. It is highly toxic to domestic animals. Malathion, upon repeated prolonged careless use, may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid. Atropine is antidote.

Contents:

MAL-MAR

GRAIN PROTECTANT

AND

BIN SPRAY

*Direct Application
to grain in bulk or
as a top dressing.*

Contains 5 Lbs. Premium Grade Malathion Per Gal.

ACTIVE INGREDIENTS:

Malathion (0, 0-dimethyl dithiophosphate of diethylmercaptosuccinate)	57%
Aromatic petroleum hydrocarbons	33%
INERT INGREDIENTS	10%

**CAUTION: Keep out of reach of children.
See side panel for additional cautions.**



DIRECTIONS FOR USE

CAUTION:
PROTECTION OF STORED
as wheat, oats, rice, corn, rye,
barghum, , and field or
against confused flour beetle, rice
weevil, saw-toothed grain
n beetle, red flour beetle, rusty
esser grain borer, and Indian
ly 1 pint of MAL-MAR in 2 to
ter per 1000 bushels. Apply as
ng loaded or turned into final

ED GARDEN SEEDS can be
st the above grain pests with
pint MAL-MAR in 1 to 2 1/2
r per 500 bushels of seed.

**Do not use for food, feed
purposes.**

CAUTION:
PROTECT STORED GRAINS from
n meal moth, Angoumois grain
MAL-MAR to the surface of clean
grain at the rate of 1/2 pint in
of water per 1000 square feet
e area. Apply the spray evenly
ce of the grain. Apply imme-
rain is loaded into storage and
sary.

U.S.A. Reg. No. 1307-55

DIRECTIONS FOR USE

3. RESIDUAL BIN SPRAY:

FOR A RESIDUAL WALL, F
MACHINERY SPRAY IN GRA
TORS, in treating truck beds, E
ships' holds before loading gr
gallon of MAL-MAR per 25 gall
making thorough application. Be
spray, clean elevators, box cars, etc
Remove and burn all sweepings a

CAUTION: Harmful if swa
absorbed through the skin. Avo
breathing of spray mist. Avoid
skin, eyes, or clothing. Wash tho
soap and water after using. Avoi
tion of feed or foodstuffs. There
application where milk is stored
or in dwellings. Keep out of reac
and domestic animals. Malathion, u
or prolonged careless use, may c
terase inhibition. If pois
get prompt medical aid. Atropine

Contents: