

CAUTION
Harmful if s

AVOID breathing vapors. AVOID contact with skin. AVOID contact with feed and foodstuffs. AVOID contact with water. Keep out of reach of children.
Note to Physicians: Malathion may cause chills and fever.
This product is toxic to fish. Keep out of lake and stream. Do not apply near mosquito larvae control work. Do not apply near water when weather conditions favor drift. Rinse equipment with water before cleaning. Clean equipment only as specified on this label.

DIRECTIONS FOR FLY CONTROL: For use in and around animal quarters, around yards, around homes, and around

STRAIGHT MALATHION SPRAYS

Amount Spray	Amount Malathion Emulsifiable Insecticide
2-1/2 gals.	1 lb.
12 gals.	5 lbs.
100 gals.	40 lbs.

Apply the spray at the rate of one gallon per acre and two gallons per 1,000 square feet of light or congested areas.
FLEAS ON DOGS AND CATS: Direct Application of Malathion 57% Emulsifiable Insecticide, per 100 square feet of area, using a pump sprayer. Repeat treatment in three weeks.
Animal Quarters: For control of fleas in kennels, dog houses, and cat houses, use Malathion 57% Emulsifiable Insecticide at the rate of 1 gallon per 1,000 square feet of area. Repeat treatment in three weeks.
FOR MOSQUITO CONTROL: Mosquito larvae and pupae are highly susceptible to Malathion. Mix with water at the rate of 1 gallon per 1,000 square feet of area. Repeat treatment in three weeks.
FOR HOUSEHOLD INSECTS: Roaches, Ants, and other household insects can be controlled by spraying in and about homes, and in animal quarters. Use Malathion 57% Emulsifiable Insecticide at the rate of 1 gallon per 1,000 square feet of area. Repeat treatment in three weeks.


VEGETABLES
DOSAGE PER ACRE DAY TO HARVEST RESIDUE TOLERANCE

TOMATOES
BEANS
POTATOES
CABBAGE
EGG PLANT
LETTUCE
SPINACH
ALFALFA
CLOVER
TOBACCO

FIELD CROPS

FRUITS
AMOUNT PER GALLONS DAYS TO HARVEST RESIDUE TOLERANCE

APPLES
ORANGES
PEACHES
PINEAPPLES
TAPES



MAGNOLIA 57% EMULSIFIABLE MALATHION

ACTIVE INGREDIENTS


	By Weight
Malathion*	57.00%
Petroleum Distillate	34.96%
INERT INGREDIENTS	8.04%
	100.00%

*O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate
One gallon contains 5 pounds of Malathion

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

See Side Panel for Antidote and Additional Precautions. Do Not Reuse Container. Destroy When Empty.



E.P.A. Reg. No. 1307-112

MAGNOLIA CHEMICAL CO., INC.
2646 Rodney Lane, Dallas, Texas
P. O. Box 20179, 247-7111

ACCEPTED WITH COMMENTS

CAUTION: Harmful if swallowed

AVOID breathing of spray mist. AVOID contact with skin. Wash thoroughly. AVOID contamination of feed and foodstuffs. Keep container closed and away from heat and open flame. Keep out of reach of children and domestic animals.

Note to Physicians: Malathion may cause cholinesterase inhibition. Atropine is antidotal.

This product is toxic to fish. Keep out of lakes, streams or ponds. (Except for mosquito larvae control work). Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

FOR FLY CONTROL. For use in and around buildings which house domestic animals, around yards, around homes, and around meat processing establishments.

STRAIGHT MALATHION SPRAYS

Amount Spray	Amount Malathion Emulsifiable Insecticide
2-1 2 gals.	1 cup
12 gals.	1 quart
100 gals.	2 ga s.

BAIT SPRAYS (WITH SUGAR)

ADD		Molasses or Corn Sugar
Syrup	or	1 cup
2-1 2 lbs.		1 quart
20 lbs.		2 ga s.

Apply the spray at the rate of one gallon per 1,000 square feet on painted surfaces and two gallons per 1,000 square feet on unpainted surfaces where flies alight or congregate. Use Unsulfurized Molasses

FLEAS ON DOGS AND CATS. Direct Application - Apply spray of 1 oz. of Malathion 57% Emulsifiable Insecticide, per 1 gallon of water. Wet the animal thoroughly using a hand sprayer. Repeat treatments may be needed within two or three weeks.


Animal Quarters - For control of fleas in kennels, pens, yards, lairs and other houses. Use spray of 5 oz. of Malathion 57% Emulsifiable Insecticide per gallon of water at rate of 1 gallon per 1,000 square feet. Use garden sprayer. Remove manure and debris before spraying. Repeat treatment three to five weeks if required.

FOR MOSQUITO CONTROL - Mosquito larvae in standing water. Apply Malathion 57% Emulsifiable Insecticide at rate of 13 fluid ounces (approximately 1 1/2 pounds active Malathion) per acre. Mix in sufficient water to give fairly even coverage when applied by air or ground equipment.

FOR HOUSEHOLD INSECTS - Roaches, Ants, Spiders, Silverfish, Crickets, Clover-wites, Earwig, etc. in and about homes, dairies and food processing plants. To make water spray, add 3 tablespoons of Malathion 57% Emulsifiable Insecticide to 1 quart of water (1 1/2 ounces per gallon). These water-based sprays may stain fabric surfaces. Apply by means of coarse spray nozzle to walls, floors, baseboards, trim boards, under sinks, stove, etc. to the areas frequented by insects.

BEETLES - In dwellings. Use Malathion 57% Emulsifiable Insecticide at rate of 2 to 4 tablespoons per gallon of dechlorinated water. Apply directly to mattress surfaces in sufficient quantity to "soak" the fabric and underneath the bed and underneath the bedspread to reach all possible hiding places.

Net Contents



MAGNOLIA 57% EMULSIFIABLE MALATHION

ACTIVE INGREDIENTS

	By Weight
Malathion*	57.00%
Petroleum Distillate	34.96%
INERT INGREDIENTS	8.04%
	<u>100.00%</u>

*O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate

One gallon contains 5 pounds of Malathion


CAUTION

KEEP OUT OF THE REACH OF CHILDREN

See Side Panel for Antidote and Additional Precautions. Do Not Reuse Container. Destroy When Empty.

E.P.A. Reg. No. 1307-112

MAGNOLIA CHEMICAL CO., INC.
2646 Rodney Lane, Dallas, Texas
P. O. Box 20179, 247-7111



ACCEPTED WITH COMMENTS

