

FOR USE AS
SANITIZER

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR FERTILIZING POISON REGISTERED
UNDER NO. 1317-48

Apr. 27, 1972

Comments

TROUBLE SHOOTER

CLEANER-SANITIZER

Use 4 ozs. av. TROUBLE SHOOTER to each 2-1/2 gallons of water to make an efficient sanitizer, which carries about 200 parts per million available chlorine. This should be left in contact with the work for a full 2 minutes.

NOTE: Half the above concentration (100 ppm) may be used if testing facilities are available so that the solution can be adjusted periodically to prevent the concentration of chlorine from falling below 50 ppm at any time.

MANUAL CLEANING

Rinse with cool water, use 4 ounces TROUBLE SHOOTER to 5 gallons of hot water. Scrub all surfaces. Rinse and air dry.

A Chlorine Bearing Detergent

HOLDS DOWN BACTERIA COUNT

CLEANS FASTER AND EASIER

HIGH SUDS FOR HARD OR SOFT WATER

for use on

Farm Tanks, Milking Machines, Utensils, and Most All Equipment

Around

DAIRIES AND CREAMERIES

Active Ingredients:

Sodium Hypochlorite	1.68%
Sodium Linear Dodecyl Benzene Sulfonate	4.17%
Sodium Phosphate as Na ₂ PO ₄ • 12H ₂ O	47.70%
Inert Ingredients	46.45%
Total	100.00%

E.P.A. Reg. No. 1317-48-AA

AN-FO MANUFACTURING CO.

OAKLAND, CALIFORNIA 94601

WARNING! KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANEL FOR ADDITIONAL WARNINGS

SPECIAL CLEANING TREATMENT FOR FARM TANKS

After tank is emptied, rinse with cold water. Apply TROUBLE SHOOTER to sides and bottom of wet tank, using 4 or 5 ounces to a 500-gallon tank. Scrub with wet brush. Leave on 5 minutes and rinse thoroughly with cold water.

On cold wall tanks with heavy deposit, TROUBLE SHOOTER may be used in warm water to quickly remove accumulated grease.

Scrub valves with bottle brush.

WARNING: Harmful if swallowed. Causes skin irritation. May cause eye damage. Do not get in eyes or on skin.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, get medical attention. If swallowed, drink large quantities of milk, egg whites or gelatin solution. Call a physician.

Do not re-use empty container—destroy it by burying with waste or burning. Stay away from smoke or fumes.

Avoid contamination of foods