

# FLY-DU INSECT KILLER

for

DAIRY • HOUSE • CREAMERY  
BAKERY • HOSPITAL • RESTAURANT  
FEED STORE • FOOD INDUSTRY

Kills . . .

<u>FLIES</u>	<u>ROACHES</u>
<u>ANTS</u>	<u>SILVERFISH</u>
<u>MOTHS</u>	<u>FRUIT FLIES</u>
<u>FLEAS</u>	<u>MOSQUITOES</u>


Net Wt.  
14 oz.

**CAUTION:** KEEP OUT OF REACH OF CHILDREN  
CONTENTS UNDER PRESSURE  
SEE BACK PANEL FOR OTHER PRECAUTIONS

### INGREDIENT STATEMENT

ACTIVE INGREDIENTS	
Pyrethrins .....	0.30%
Piperonyl Butoxide Technical* .....	1.50%
Petroleum Hydrocarbons .....	18.20%
INERT INGREDIENTS .....	80.00%
	100.00%

\*Equivalent to 1.20% of (butylcarbityl) (6-propylpiperonyl) ether and 0.30% of related compounds.



### DIRECTIONS FOR USE

**TO OPERATE**—Remove protective cap. Hold can upright and point hole in button toward area or object to be sprayed. Depress button to release mist. Hold dispenser two (2) feet from any surface when spraying.

**INDOOR SPRAYING**—For **FLIES, GNATS** and **MOSQUITOES**: Shut doors and windows and spray toward ceiling at the rate of three (3) to five (5) seconds per each 1000 cubic feet of space (10' x 10' x 10'). For best results keep room closed for at least fifteen (15) minutes after spraying. DO NOT remain in treated areas and ventilate the area after treatment is completed.

**ROACHES & SILVERFISH**—To kill roaches and silverfish direct spray into cracks, crevices and other secluded places. Apply liberally. Spray at insects when they appear. Repeat treatment as necessary.

**ANTS**—To kill ants spray directly at insects as well as ant trails and other places where ants gather. Repeat treatment as necessary.

**FLEAS**—To kill fleas spray infested quarters liberally. Apply directly to rugs, floors, baseboards and animal's sleeping quarters. Repeat treatment as necessary.

**MOTHS**—Remove all clothing from closet and spray all surfaces thoroughly. Close doors for 30 minutes. Spray surfaces of garments including seams, folds and lapels before returning to closet. Repeat treatment every thirty (30) days if garments are not stored in moth tight containers.

**FOOD PROCESSING PLANTS**—Food should be removed or covered during treatment. All food processing surfaces should be covered during treating or thoroughly cleaned before use. When using product in these areas, apply only when plant is not in operation.

**DAIRY BARN AND MILK ROOMS**—To control flies in dairy barns and milk rooms direct spray toward the ceiling. Release mist at the rate of three (3) to five (5) seconds per each 1000 cubic feet of space. Release sufficient material to get a thin mist of insecticide in the air. For best results keep all doors and windows closed when possible. Repeat treatment as necessary.

**CAUTION**—Cover fishbowls when spraying. Remove pets from premises. To prevent contamination of milk and utensils, keep covered. Do not contaminate food or feed stuffs. Keep out of reach of children.

**EXTREMELY FLAMMABLE: CONTENTS UNDER PRESSURE. KEEP AWAY FROM FIRE, SPARKS, AND HEATED SURFACES. DO NOT PUNCTURE OR INCINERATE CONTAINER. EXPOSURE TO TEMPERATURES ABOVE 130° F. MAY CAUSE BURSTING.**

EPA Reg. No. 1317-74-AA

EPA Est. 27417-CA-1

Packed for  
**AN-FO MANUFACTURING COMPANY**  
Oakland, California 94601

**BEST AVAILABLE COPY**