

ACCEPTED

JUL 12 1974

UNDER FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. *1327-28* SUBJECT
TO ATTACHED COMMENTS.



FUMIGATOR
No. 20 M

This fumigator will fumigate 20,000 cu. ft.
for use in greenhouses only

SEE CAUTION ON REAR PANEL

Active Ingredients:
*Gamma Isomer of Benzene Hexachloride - 11.2%
Inert Ingredients: 88.8% *From Lindane

CAUTION!

KEEP OUT OF REACH OF CHILDREN
(See back panel for additional cautions.)

For use against greenhouse aphids, grasshoppers,
adult white flies and adult leaf tiers.

Net Weight — 5.3 oz.

Manufactured by
FULLER SYSTEM, INC., WOBURN, MASS., U.S.A.

DIRECTIONS

Shake contents before using. Unscrew top cover of container as directed when ready to use.

Place fumigators in single row on the ground in a central walk. Space them evenly for proper distribution of smoke.

Close all ventilators, doors and other openings. Starting with fumigator at farthest end of greenhouse away from exit, light sparkler at handle end and insert into top half inch of contents. Proceed to next fumigator, etc., until last fumigator is ignited, then leave greenhouse which should remain closed for at least four hours. Do not resume work in greenhouse until thoroughly ventilated. Cut wide open blooms before fumigating.

Ventilate gradually, preferably within eight hours — particularly when crops are in bloom or when growth is unusually soft.

Cloudy day, late afternoon or night fumigations are recommended, at temperatures between 60 - 80° F. Avoid fumigating when windy or in the presence of excessive moisture.

Repeat fumigation when necessary if reinfestation occurs.

CAUTION

Avoid contact with skin. Avoid inhalation of vapors. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans.

Do not remain in greenhouse during fumigation without wearing full-face mask of a type passed by the U.S. Bureau of Mines for Lindane protection and protective clothing to minimize the danger of absorption of the vaporized material through the skin. May be absorbed through the skin. In case of contact, wash immediately with soap and water.

NON-WARRANTY

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

CAUTION

DO NOT USE ON EDIBLE FOOD CROPS EXCEPT AS FOLLOWS:

- TOMATOES:** Allow 1 day between treatment and harvest.
- CUCUMBERS:** Allow 1 day between treatment and harvest.
- LETTUCE:** Allow 5 days between treatment and harvest.

E.P.A. Reg. No. 1327-28

DO NOT OPERATE THIS CAN NEAR FLAMMABLE MATERIAL

