

1339-6

ACCEPTED

Feb-27-1969

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT

ED UNDER NO. 1339-6 SUBJECT TO ATTACHED COMMENTS.

COTTON STATES TOXAPHENE 20% DUST

DIRECTIONS FOR USE

USDA Reg. No. 1339-6

DIRECTIONS FOR USE

Cotton States 20-0 Toxaphene Dust is formulated primarily for use in controlling insects attacking many crops. This material can be effectively applied with conventional hand or power dusters, or by airplanes.

SMALL GRAINS: For control of fall armyworms, yellow striped armyworms, grasshopper, chinch bug, climbing cutworms and surface feeding cutworms in rice, wheat, oats, barley and rye, apply Cotton States 20-0 dust at a rate of 10 lbs. per acre. More than one application may be necessary for control throughout the season.

Do not apply to small grains more than once after heads have begun to form. Do not apply within 7 days of harvest on oats, wheat and rye. Do not apply within 14 days of harvest on rice and barley.

COTTON: For control of thrip, fleahopper, rapid plant bug and tarnished plant bug, apply to young growing cotton at the rate of 7 to 8 lbs. per acre when insects appear. Repeat as necessary for control. For control of boll weevils, cotton bollworms and cotton leafworms, apply Cotton States 20-0 dust at rates of 10 to 20 lbs. per acre when insects appear and repeat every 5 days until insects are under control.

Thorough uniform coverage is essential to efficient control of insects.

CUTWORMS: For control of cutworms in peppers, beans, eggplant, lettuce, melons, and tomatoes, apply to soil surface around base of plant at the rate of 10 lbs. per acre. For control of cutworms in broccoli, brussel sprouts, and cauliflower, apply to soil surface around base of plants at the rate of 12 to 15 lbs. per acre when insects appear. For control of cutworms in strawberries, apply to soil surface around base of plants at rate of 15 lbs. per acre. Make no application after appearance of foliage to be eaten or fed to dairy animals or animals being finished for slaughter. Make no application after edible parts begin to form.

TOMATOES: For control of tomato fruitworm, tomato pinworm, armyworms and climbing cutworms, apply 25 lbs. per acre as needed when insects appear when fruit of main crop is setting. Repeat at two week intervals if necessary. For control of blister beetles, apply 25 lbs. per acre. For flea beetles and hornworms, apply 15 to 20 lbs. per acre as needed. Uniform coverage is necessary for best results.

Make no application on tomatoes within 5 days of harvest. Remove excessive residues by washing and rinsing.

ACTIVE INGREDIENTS:

*Toxaphene	20.00%
INERT INGREDIENTS ...	80.00%
TOTAL	100.00%

*Technical Chlorinated Camphene (Chlorine Content 87% to 89%)

Broccoli, Brussels Sprouts, Cauliflower and lettuce, For control of caterpillars in above crops, apply 20 lbs. per acre. Make no application to the above after appearance of plant parts to be eaten or marketed.
STRAWBERRY: For control of crown borer and spittle bugs, apply 15 to 20 lbs. per acre as needed when insects appear. Make no application to strawberries while fruit is on the plant.

CAUTION

Keep Out of Reach of Children

May be fatal if swallowed. DO NOT breathe dust. Do not get on skin, in eyes or on clothing. Wash thoroughly. Do not contaminate food or feed-stuffs. Keep away from children and domestic animals (particularly dogs). To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material. DO NOT use for anything other than specified on this label, and at recommended dosages. Do not pasture dairy animals or animals being fattened for slaughter in treated field. Do not apply nor allow to drift where livestock are grazing. Plants treated with this material should not be fed nor should ensilage from such plants be fed to dairy animals or animals being finished for slaughter. This product is poisonous to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on the label. Do not apply when weather conditions favor drift of dust from crop areas treated.

ANTIDOTE

Internal—Give tablespoonful of salt in glass of warm water and repeat until vomit fluid is clear. CALL A PHYSICIAN.
External—Wash thoroughly with soap and water immediately.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Manufactured By



MONROE, LOUISIANA

1339-6

ACCEPTED
Feb-27-19
UNDER THE FEDERAL INSECT
POISONIDE AND RODENTICIDE

ED UNDER NO. 1339-6
TENTATIVE COMMENTS.

COTTON STATES TOXAPHENE 20% DUST

DIRECTIONS FOR USE

Cotton States 20-0 Toxaphene Dust is formulated primarily for use in controlling insects attacking many crops. This material can be effectively applied with conventional hand or power dusters, or by airplanes.

SMALL GRAINS: For control of fall armyworms, yellow striped armyworms, grasshopper, chinch bug, climbing cutworms and surface feeding cutworms in rice, wheat, oats, barley and rye, apply Cotton States 20-0 dust at a rate of 10 lbs. per acre. More than one application may be necessary for control throughout the season.

Do not apply to small grains more than once after heads have begun to form. Do not apply within 7 days of harvest on oats, wheat and rye. Do not apply within 14 days of harvest on rice and barley.

COTTON: For control of thrip, fleahopper, rapid plant bug and tarnished plant bug, apply to young growing cotton at the rate of 7 to 8 lbs. per acre when insects appear. Repeat as necessary for control. For control of boll weevils, cotton bollworms and cotton leafworms, apply Cotton States 20-0 dust at rates of 10 to 20 lbs. per acre when insects appear and repeat every 5 days until insects are under control.

Thorough uniform coverage is essential to efficient control of insects.

CUTWORMS: For control of cutworms in peppers, beans, eggplant, lettuce, tomatoes and tomatoes, apply to soil surface around base of plant at the rate of 10 lbs. per acre. For control of cutworms in broccoli, brussel sprouts, and cauliflower, apply to soil surface around base of plants at the rate of 12 to 15 lbs. per acre when insects appear. For control of cutworms in strawberries, apply to soil surface around base of plants at rate of 15 lbs. per acre. Make no application after appearance of foliage to be eaten or fed to dairy animals or animals being finished for slaughter. Make no application after edible parts begin to form.

TOMATOES: For control of tomato fruitworm, tomato pinworm, armyworms and climbing cutworms, apply 25 lbs. per acre as needed when insects appear when fruit of main crop is setting. Repeat at two week intervals if necessary. For control of blister beetles, apply 25 lbs. per acre. For flea beetles and hornworms, apply 15 to 20 lbs. per acre as needed. Uniform coverage is necessary for best results.

Make no application on tomatoes within 5 days of harvest. Remove excessive residues by washing and rinsing.

USDA Reg. No. 1339-6

DIRECTIONS FOR USE

Broccoli, Brussels Sprouts, Cauliflower and lettuce. For control of cutworms in above crops, apply 20 lbs. per acre. Make no application to above after appearance of plant parts to be eaten or marketed.
STRAWBERRY: For control of crown borer and spittle bugs, apply 15 to 20 lbs. per acre as needed when insects appear. Make no application to strawberries while fruit is on the plant.

ACTIVE INGREDIENTS:

*Toxaphene	20.00%
INERT INGREDIENTS	80.00%
TOTAL	100.00%

*Technical Chlorinated Camphene (Chlorine Content 67% to 69%)

CAUTION

Keep Out of Reach of Children

May be fatal if swallowed. DO NOT breathe dust. Do not get on skin, eyes or on clothing. Wash thoroughly. Do not contaminate food or feed stuffs. Keep away from children and domestic animals (particularly dogs). To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material. DO NOT use for anything other than specified on this label, and at commended dosages. Do not pasture dairy animals or animals being fattened for slaughter in treated field. Do not apply nor allow to drift where livestock are grazing. Plants treated with this material should not be used for ensilage from such plants be fed to dairy animals or animals being finished for slaughter. This product is poisonous to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply in any manner specified on the label. Do not apply when weather conditions favor drift of dust from crop areas treated.

ANTIDOTE

Internal—Give tablespoonful of salt in glass of warm water and repeat until vomit fluid is clear. CALL A PHYSICIAN
External—Wash thoroughly with soap and water immediately.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of such use.

Manufactured By



MONROE, LOUISIANA