

TESTED

9-24-71
1339-68

UNDER THE FEDERAL INSECTICIDE ACT
FOR THE FEDERAL REGISTER.
EFFECTIVE DATE OF REGISTER.
IN ATTACHED COMMENTS

COTTON STATES MALATHION EMULSIFIABLE 57

U. S. D. A. Reg. No. 1339-68

CAUTION

Keep Out Of Reach Of Children

HARMFUL if swallowed, inhaled or absorbed through the skin.

AVOID breathing spray mist.

AVOID contact with skin or eyes. Bathe promptly with soap and water after working with Malathion. Wash clothing before re-use.

AVOID contamination of feed and foodstuffs.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Note to physician: Malathion may cause cholinesterase inhibition. Atropine is antidotal.

This product will kill fish. Keep out of any body of water.

Do not apply when weather conditions favor drift from areas treated: Do not apply where runoff is likely to occur; Do not contaminate water by cleaning of equipment or disposal of wastes; Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchant ability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

ACTIVE INGREDIENTS:

*Malathion 57%

Aromatic Petroleum Distillates 35%

INERT INGREDIENTS: 8%

TOTAL 100%

*Technical 95%

*O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate

CONTAINS 5 LBS. TECHNICAL MALATHION PER GALLON

Directions for Use—

For aerial application, apply recommended rate in a minimum of 3 gal. water per acre. For ground application, apply in sufficient water for uniform coverage.

Crop	Pest	Amount to use per acre	Days between last applic. and harvest or grazing
Pasture and Range grass	Armyworms Grasshoppers	2 pints	0
Tomatoes (outdoor)	Spider mites Aphids Drosophila	1½ pint 1 pint 2½ pints	1 1 1
Peas	Pea Aphid Pea Weevil	1½ pints 1½-2 pints	3 3
Beans	Mexican Bean Beetle Spider Mites Japanese Beetle Aphids	1½ pints 1-1½ pints 1½-2 pints 2 pints	1 1 1 1
Broccoli Collards Cabbage Turnips Mustard Greens	Aphids	1½-2 pints	7
Cucumbers Melons	Aphids Spider Mites Pickleworm	1½ pints	1

Crop	Pest	Amount to mix with 100 gal. water	Special Instructions
Ornamentals	Lace Bugs Spider Mites Whitefly Thrips Tarnished Plant Bug Euonymus Scale Mealybugs Azalea Scale Tent Caterpillar Bagworms	2 pints	Spray to run off. Injury may occur on ferns, hickory, viburnum, lantana, crassula and canaertii juniper. Boston, pteris, and maidenhair ferns, petunias, spirea, white pine and maples are also sensitive.
Cattle (beef and non-milking) Horses	Lice Ticks Hornfly	1-2 gal.	Treat animals thoroughly. Repeat applications at 10-14 day intervals until pest is controlled. Do not treat non-milking cattle within 2 weeks of freshening.
Hogs	Lice Sarcoptic Mange	1 gal.	Cover all body surfaces including the inside of the ears. A second treatment may be necessary in 10 days. Hogs should be kept out of sun and wind for a few hours after treatment.

Manufactured by



Monroe, Louisiana