


PARATHION-8 E.

Contents U. S. Gallons

(8 lbs. Parathion per gal.)



POISON



ACTIVE INGREDIENTS

Parathion (OO methyl 2-p nitrophenyl methyl phosphorothioate)

Xylene Range Aromatic Solvent

INERT INGREDIENTS

TOTAL

100

RECOMMENDATIONS

BEANS Aphids, Armyworms, Curculionid Beetles, Leafhoppers, Leaf Miners, Leaf Roller, Lima Pod Borer, Mexican Bean Beetle, Stinkbugs, Thrips, apply 1/2 pt per acre. Repeat applications at 7 to 10 day intervals and continue TO NOT APPLY WITHIN 7 days of harvest.

COLE CROPS Cabbage, Broccoli, Brussels Sprouts, Cauliflower, Kale, Mustard, Turnip, Aphids, Armyworms, Cabbage Looper and Imported Cabbageworm, Leaf Miners, Thrips, apply 1/2 pt per acre. DO NOT APPLY ON TURNIPS WITHIN 14 days of harvest, 21 days if the insecticide is used on a leaf on other crops within 14 days of harvest.

COTTON Cotton, Aphid, Cotton Leafworm, Fleahoppers, Spider Mites, Stinkbugs, apply 1/2 pt per acre. Cotton Leaf Perforator, apply 1/2 pt per acre. DO NOT APPLY WITHIN 7 days of harvest.

CUCURBITS Cucumber, Squash, Melons, Aphids, Cucumber Beetle, Melonworm, Pickleworm, Leaf Miners, Squash Bug and Burrowing in watermelons, apply 1/2 pt per acre. DO NOT APPLY UNLESS PLANTS ARE DRY. DO NOT APPLY BEFORE VINING OR WITHIN 14 days of harvest on cucumbers and squash, 21 days of harvest on melons.

FOOD CROPS Corn, Broomrape, Clover, Oats, Rye, Wheat, Aphids, Armyworms, Blister Beetles, Grasshoppers, apply 1/2 pt per acre at first sign of infestation. DO NOT APPLY WITHIN 15 days of harvest or 30 days if used on seed.

SOYBEANS Aphids, Leaf Miners, Red Spider Mites, Thrips, apply 1/2 pt per acre. DO NOT APPLY WITHIN 7 days of harvest.

CELERY Aphids, Leaf Miners, Stinkbugs, apply 1/2 pt per acre. DO NOT APPLY WITHIN 7 days of harvest.

LETTUCE Aphids, Beet Leafhopper, Beetle, Leafhopper, Flea Beetle, apply 1/2 pt per acre. DO NOT APPLY WITHIN 7 days of harvest. Do not use on lettuce 14 days of harvest on leaf and root lettuce.

PUMPKIN Aphids, Flea Beetle, Leaf Miners, Thrips, apply 1/2 pt per acre. DO NOT APPLY WITHIN 7 days of harvest.

POTATOS Aphids, Armyworms, Beet Leafhopper, Beetle, Flea Beetle, Leafhoppers, Leafminer, Plant Bug, Leaf Miner, Green Stinkbugs, apply 1/2 pt per acre. DO NOT APPLY WITHIN 7 days of harvest.

TOMATO Aphids, Beet Leafhopper, Beetle, Flea Beetle, Stinkbugs, apply 1/2 pt per acre. Flea Beetle, apply 1/2 pt per acre. DO NOT APPLY WITHIN 7 days of harvest.

STRAWBERRIES Flea Beetles, Flea Cicadas, Flower Weevil, Leaf Miner, Root Borer, apply 1/2 to 1 pt per acre. Red Spider Mite, apply 1/2 pt per acre. DO NOT APPLY WITHIN 14 days of harvest.

GENERAL DIRECTIONS

Mix in a spray mixture as follows:

Add about half the total amount of water to be used to the spray tank.

Start agitation and pump system and add proper amount of Parathion-8 E.

Continue mixing until all Parathion-8 E. is dissolved.

Directions for Use:

Parathion-8 E. may be applied by either hand equipment or power equipment. For power equipment, dilute in sufficient water to give a concentration of 8 lbs. Parathion-8 E. per 100 gallons of water per acre.

Application should be made only by trained persons using appropriate water equipment. DO NOT APPLY WITH HAND EQUIPMENT.

If mixture does not mix readily, it is best to separate as a dry powder and mix as many pounds as possible of the mixture with water in a power sprayer. If the mixture does not mix readily, add little water at a time. ALWAYS dilute with water before use. Do not use water which is hard or contains lime.

Reserve an adequate amount of the mixture for use in case of an emergency that pest control residues and the amount of water applied may vary. Do not use a mixture prepared by the Federal Government. Use only a few minutes before use. Repeat applications at 7 to 10 day intervals and continue to apply up to 7 days before harvest. Do not use a mixture of Parathion-8 E. and other insecticides or fungicides. THE GROWER IS RESPONSIBLE for residues and the amount of water applied. Do not use a mixture of Parathion-8 E. and other insecticides or fungicides.

U. S. PATENT OFFICE

Manufactured by

Cotton States Chemical Co., Inc.

Monroe, Louisiana