

Cotton States[®] 10% Sevin[®] Dust

NET
WT. _____

GENERAL DIRECTIONS

Cotton States 10% Sevin Dust is formulated for control of many insects attacking cotton, beans and potatoes. It can be applied in conventional hand or power dusters or by airplane. Uniform, thorough coverage is necessary for efficient control.

SUGGESTIONS FOR USE

ON COTTON: For control of fleahoppers and Thrips, apply 5 to 10 lbs. per acre.

For control of Boll Weevils, Cotton Bollworms, Pink Bollworms, Cotton Leafworms, Lycus Bugs and Cotton Leaf Perforator, apply Cotton States 10% Sevin Dust at the rate of 15 to 25 lbs. per acre depending on rankness of foliage and degree of insect infestation.

Begin all applications when insects appear and repeat every 5 to 7 days until insects are under control. When used on a regular schedule at above rates Cotton Aphids will be suppressed. Do not graze treated fields or feed trash to livestock.

ON BEANS AND POTATOES: For control of Mexican Bean Beetle on beans and the Colorado Potato Beetle on potatoes, apply at the rate of 10 lbs. per acre.

For control of Leafhoppers on beans and Leafhoppers and Flea Beetles on potatoes, apply at the rate of 10 to 15 lbs. per acre depending on the degree of infestation and rankness of foliage.

For control of Lycus Bugs on beans and potatoes, apply 10% Sevin Dust at the rate of 20 lbs. per acre.

For control of the above insects on beans and potatoes, begin applications when insects appear, and repeat at 7 to 10 day intervals until insects are controlled. Thorough, uniform foliage coverage is essential to efficient insect control.

Do not graze treated fields or feed treated vines to dairy animals or animals being finished for slaughter.

CAUTION

Keep Out of Reach of Children

Harmful if swallowed or inhaled. Do not breathe dust. Do not take internally. Avoid breathing of dust. **SKIN CONTACT MAY BE HARMFUL.** Wash hands before eating. Take shower or bath after work. **WEAR REGULAR LONG SLEEVED WORK CLOTHING.** Change to clean clothing daily. Avoid prolonged or repeated contact with the skin, wash thoroughly after using. Avoid storage near feed or food products. **NOTE FOR PHYSICIAN:** (Carbaryl is a moderate, reversible, cholinesterase inhibitor.) Atropine is Antidotal. For protection of bees, avoid use, if possible, during periods when honey bees are visiting the crops. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

To protect fish and wildlife, do not contaminate streams, lakes, or ponds with this material. This product is poisonous to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be harmed.

Keep container tightly closed when not in use.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

EPA Reg. No. 1339-136

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-Methylcarbamate) 10.0%

INERT INGREDIENTS 90.0%

TOTAL 100.0%

® Trademark of Union Carbide Corp. for Carbaryl—
(1-Naphthyl N-Methylcarbamate)

DO NOT REUSE CONTAINER

DESTROY WHEN EMPTY

Manufactured by



MONROE, LOUISIANA

ACCEPTED

Jul 17, 1972