

"RS" RESIDUAL INSECTICIDE

ACTIVE INGREDIENTS	85%
* o-Isopropoxyphenyl methylcarbamate	1.00%
Pyrethrins	03%
Piperonyl Butoxide	2369
Odorless base oil	83.7345
INFRT INGREDIENTS	
Isopropyl alcohol	9.987
Methylene Chloride	5.00%
Essential oil	.02%
- #U.S. PAT.# 3,111,530 & CAN . TAN PAT. # 7	25 ,20 4

85 123 074, insectionse is excellent for use in Mills, Bakeries, Fond Markets, Furniture Storage and other Warehouses, Hospitals, and Institutions. When used as directed R3 is effective in killing resistant roaches, water-bugs suverfish, carpet beetles, Jidd 13 1111 ing insects

Die Thicker De hist

* *** · (1

the second sign, and idvertible r de la companya de

outdoors

CAUTION KEEP OUT OF REACH OF CHILDREN—HARMFUL IF SWALLOWED AVOID CONTACT WITH SKIN OR F 5. MAY BE ABSORBED THRU SKIN. Use in food processing plants only when plant is not in operation. Avoid contact with eyes, skin and clething. Use with adequate ventilation. Do not permit children to contact treated surfaces until dry, Do not reuse containers. Destroy containers by perforating or crushing. Bury or discard in a safe place.

the control of the second second second second A district on figure and or agree there are a residence of contamination of find to deriff in first in a Dinner star

In case of illness, get prompt medical aid. Atropine sulfate is antidotal. And the second of the second

S SAINER

IS-WEIL MFG. CO., INC. ::

21 - Sign To C

2714 35th ST TAMPA FLA

A CLEANER FOR EVERY NEED:

Cotton States Sevin Super Tomato Dust

Manufactured By




MONROE, LOUISIANA

COTTON STATES SEVIN SUPER TOMATO DUST is a combinafon insecticide, fungicide dust formulated for use on Tomatoes. When applied in accordance with directions, this Blights, Southern Blights, Anthracnose and Septoria Leaf Spot well as the following insects. Tomato ruitworm, Tomato flornworms, Fall Armyworms, Flex Hopper, European Corn Borer, Stink Buds, Striped and Septoria Leaf Spot Hopper, European Corn Borer, Stink Buds, Striped and Striped duster or by airplane.

DIRECTIONS FOR USE

ON TOMATOES: For control of the above named insects and diseases apply Cotton States Super Tomoto dust at the rate of 20 to 40 lbs. per acre depending on size of plant and rankness of foliage. Begin applications when fruit on first cluster begin to form. Repeat applications every seven days. Repeat immediately any application washed off by rain. Complete and uniform coverage is necessary for effective control.



CAUTION

DO NOT apply this dust within 1 day of harvest. Tomatoes should be processed to remove residues before marketing.

WARNING

KEEP OUT OF THE REACH OF CHILDREN

Harmful if swallowed or inhaled. Do not take internally Avoid breathing dust. Avoid contact with skin and eyes. For skin contact, wash with plenty of soap and water. For eyes, flush with water for 15 minutes. Get prompt medical attention. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs.

Seller warrants that the product conforms to its chemica. description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any ther warranty of nerchantability or fitness for a particular surpose, expressed or implied, extends to the use of this product contrary to lake) instructions, or under abnormal conditions or under conditions not reasonably torsecable to while, and buyer assumes the risk of any such ~ -1

ACTIVE INGREDIENTS:

Zinc Ethylene Bisdithiocarbamate (Zineb)	6.0%
Carbaryl (1-naphthyl N-methylcarbamate)	10.0%
INERT INGREDIENTS:	84.0%
TOTAL	100.0°

*Trademark of Union Carbide Corp. for the active ingredients. Carbaryl (1-naphthyl N-methylcarbamate).