

Cotton States 5% Sevin® Garden Dust

U. S. D. Reg. No. 1339-159

DIRECTIONS

APPLICATION: Begin applications when insects or damage is first observed and repeat on a regular 7 to 10-day schedule or as necessary for good insect control.

VEGETABLES

BEANS: To control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles, and Corn Earworm—apply at the rate of 20 pounds per acre or 1 ounce per 50 feet of row. Where Mexican Bean Beetle is the only problem, a rate of 10 pounds per acre is sufficient. To control Lygus Bugs, Stink Bug and Lima Bean Pod Borer—apply 40 pounds per acre. This product can be applied up to the day of harvest.

CUCUMBER, SUMMER SQUASH: To control Pickleworm and Melonworm—apply at the rate of 10 pounds per acre. To control Cucumber Beetles, Squash Bugs, and Flea Beetles—apply at the rate of 20 pounds per acre or 1 ounce per 50 feet of row. This product can be applied up to the day of harvest.

TOMATO, EGGPLANT, PEPPER: To control Tomato Fruitworm, Fall Armyworm, Flea Beetles and Leafhoppers—apply at the rate of 20 pounds per acre or 1 ounce to 50 feet of row. For control of Tomato Hornworm, European Corn Borer and Stink Bugs—apply at the rate of 40 to 80 pounds per acre. This product can be applied up to the day of harvest.

POTATO: To control Colorado Potato Beetle—apply at the rate of 10 pounds per acre. For control of Leafhoppers and Flea Beetles apply at 20 pounds and for Lygus Bugs apply at 40 pounds per acre.

ACTIVE INGREDIENTS

Carbaryl—1-Naphthyl N-Methylcarbamate 5.0%

INERT INGREDIENTS 95.0%

TOTAL 100.0%

® Trademark of Union Carbide Corp. for Carbaryl—1-Naphthyl N-Methylcarbamate.

MANUFACTURED BY



MONROE, LOUISIANA

ORNAMENTALS

To control Apple Aphid, Bagworm, Birch Leaf Miner, Blister Beetles, Boxelder Bug, Boxwood Leaf Miner, Elm Leaf Aphids, Elm Leaf Beetle, Flea Beetles, Japanese Beetle, Lace Bugs, Leafhoppers, Leaf Rollers, Oak Leaf Miners, Orange Tortrix, Periodical Cicada, Plant Bugs, Psyllids, Rose-slug, Rose Aphids, Scale insects in the crawler stage, Tent Caterpillars, Willow Leaf Beetle and exposed Thrips—apply thoroughly to the infested plants at first sign of damage and repeat at 7-day intervals or as necessary for good insect control. To obtain scale control apply in Spring and early Summer when scale crawlers are present.

LAWNS

To control Leafhoppers, Fall Armyworm, and Sod Webworm—apply at the rate of 4 pounds per 1,000 square feet. For best results, apply after rain or watering and do not water for at least 2 days after application. Repeat in 2 to 3 weeks if necessary.

CAUTION

Keep out of Reach of Children

Harmful if swallowed or inhaled. Do not breathe dust. Do not take internally. Avoid prolonged or repeated breathing of dust. SKIN CONTACT MAY BE HARMFUL. Avoid contact. Wash hands before eating. Take shower or bath after work. WEAR REGULAR LONG SLEEVED WORK CLOTHING. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is Antidotal.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

ACCEPTED

8/6/71

UNDER THE FEDERAL INSECTICIDE,
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PEST CONTROL