

# Toxaphene-8 Emulsifiable

(8 pounds toxaphene per gallon)

## ACTIVE INGREDIENT:

Toxaphene (Technical Chlorinated Camphene containing 67-69% Chlorine)	71.0%
Aromatic Petroleum Distillate	25.0%
INERT INGREDIENTS:	4.0%
TOTAL	100.0%

## DIRECTIONS FOR USE

Cotton: Early Season—For control of thrips, fleahoppers, and plant bugs. Apply 2-1/4 to 3-3/4 pints of this material suitably diluted with water per acre. This product may be applied with either aerial or ground sprayers. With aerial application, the desired volume of insecticide concentrate should be diluted with water and applied at 3 to 4 gallons per acre. For ground applications, the desired volume of insecticide concentrate should be diluted with water and applied at 6 to 8 gallons per acre.

Make 2 to 3 applications at 7-day intervals starting at the two-leaf stage. Avoid spray drift. Do not graze dairy animals or animals being finished for slaughter on cotton fields which have been treated with toxaphene late in the season.

Wheat, Oats: For control of armyworms and cutworms, use 2-1/4 to 3 pints of this product suitably diluted and applied as indicated for cotton.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

## WARNING

### Keep Out of Reach of Children

Read Entire Label for Additional Handling Precautions.

WARNING: May be fatal if swallowed. Do not get on skin, in eyes, or on clothing. Can be absorbed through skin. Wash thoroughly after using. Do not breathe spray mist. Avoid contamination of feed or foodstuffs. Do not use on dairy animals or in dairy barns. Do not use on cucurbits.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams and ponds. Apply only as specified on the label. Rinse equipment and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

## ANTIDOTE:

Internal—Give sodium chloride 15 g. (one tablespoonful of table salt) in glass of warm water and repeat until vomit is clear. Call a doctor immediately.

External—Wash with soap and water.

1331-115  
ACCEPTED

Oct 30 1958

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND FERTILIZER ACT  
FOR COTTON AND OTHER  
ED UNDER REG. NO. 1339-195  
TO ATTACHED LABEL

Sorghum: For control of armyworms and cutworms, use 2-1/4 to 3 pints of this product suitably diluted and applied as indicated for cotton. Do not apply within 40 days of harvest. Do not apply more than once after heads start to form. Do not graze dairy animals or animals being finished for slaughter in treated fields. Do not ensile treated forage.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use.

USDA Reg. No. 1339-195

Net Contents Gal.

Manufactured by



MONROE, LOUISIANA