COTTON STATES KETOKIL NO. 63

U. S. D. A. Reg. No. 1339-206

DANGER

heep Out of Reach of Children

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. Rapidly absorbed through skin. Wear natural rubber gloves, protective clothing and goggles.

Do Not get in eyes, on skin, or on clothing. In case of contact remove clothing, wash thoroughly with soap and water, for eyes flush with plenty of water and get prompt medical attention.

DO NOT breathe spray mist or vapors. Wear a mask or respirator of a type passed by U. S. Dept. of Agriculture for Methyl Parathion protection.

KEEP ail unprotected persons and animals out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be reentered until drifting spray mist and volatile residues have dissipated. If headache, dizziness or nausea develops during or after use, get out of contaminated area at once. Get medical attention.

WEAR clean clothing daily, wash all contaminated clothing with soap and hot water before re-use.

Wash hands, arms and face thoroughly with soap and water before eating or smoking.

DO NOT contaminate feed and foodstuffs. This product is poisonous to fish and wildlife. Keep out of lakes, po ds and streams. Do not apply in any manner not specified on the label.

Eirds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.

To not feed treated forage to dairy animals or animals being finished for slaughter.

Do not apply when weather conditions favor drift of spray or dust from areas treated.

DESTROY all containers when empty, by punch ing holes in them and burying with wastes. Do not burn.

Rinse equipment and containers and dispose of wastes by burying in noncrop lands away from water supplies.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

NOT for use or storage in or around the home.

CAUTION: Do not apply within 5 days of hand picking. Workers entering fields within 24 hours after application should wear protective chollang. Do not graze daw, animals or animals being finished for slaughter in treated fields.

ACTIVE INGREDIENTS:

*Toxaphene 52.5°,

*O,O-dimethyl O-p-nitrophenyl thiophosphate 26.2°,

Xylene (Arcinatic Petroleum Distillate) 17.5°,
INERT INGREDIENTS 3.8°,

- *Technical chlorinated Camphene (Chlorine Content 67% to 69%)
- "Methyl Parathion

TOTAL

Each gallon of Cotton States Ketokil No. 63 contains 6 lbs. toxaphene plus 3 lbs. Methyl Parathion. Used as directed will control the following insects: Thrips, Fleahoppers, Plant bugs, Boll Weevils, Bollworms, Cotton Leafworms, Aphids, and Corn earworm.



POISON



100.0°_c

ANTIDOTE FOR METHYL PARATHION

IF SWALLOWED: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quite. Call a physician IMMEDIATELY. IF ON SKIN: In case of contact remove contaminated clothing and immediately wash skin with soap and water.

PHYSICIAN'S NOTE

Compound exerts muscarine-like effects i.e., stimulation of the para-sympathetic nervous system. Repeated use of a para-sympathetic depressant, such as atropine to the limit of patient's tolerance, is indicated to relieve symptoms. The administration of oxygen is helpful. Never give morphine to patient's suspected of Parathion poisoning. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Manufactured By



ACCEPTED

GENERAL DIRECTIONS

Prepare a spray mixture of Ketokil No. 63 as follows:

- 1. Add about half the total amount of water to be used to the spray tank
- 2. Start agitator or pump bypass and add proper amount of Ketokil No. 63.
- 3. Continue mixing and add remaining water.

DIRECTIONS FOR USE ON COTTON

Ketokil No. 63 may be applied by either ground equipment or aircraft. For ground equipment, dilute in sufficient water to give adequate coverage. For aerial applications, dilute in 3 to 4 gallons of water per acre.

Early Season Program: For general early season control of thrips, plant bugs and fleahoppers, use I gal. Ketokil No 63 to 12 acres Make two or three applications at 7 day intervals starting at the two-leaf stage.

Mid Season Insects. Overwintering bold weevils, plant bugs, aphids, and cotton leaf-worms are controlled with 1 gal. Ketokil No. 63 to 6 acres. Make repeat treatment at 7 day intervals as insect counts indicate they are necessary.

Late Season Control. Use one gallon to 3 acres to control boll weevils, bollworms, and aphids. Repeat applications at 5 to 7 day intervals as insect counts indicate they are necessary

The second of th