# COTION STATES KETOKIL NO. 44

USDA Reg. No. 1339 208

## **DANGER**

### heep Out of Reach of Children

POISONOUS IF SWALLOWED, INHALED OR AB SORBED THROUGH SKIN. Rapidly absorbed through skin. Vear natural rubber gloves, protective clothing and goggles.

Do Not get in eyes, on skin, or on clothing. In case of contact remove clothing, wash thoroughly with soap and water, for eyes flush with plenty f water and get prompt medical attention.

DO NOT breathe spray mist or vapors. Wear a mask or respirator of a type passed by U. S. Dept of Agriculture for Methyl Parathion protection

KEEP -II unprotected persons and animals out if operating areas or vicinity where there may be danger of drift. Vacated areas should not be reintered until drifting spray mist and volatile residues have dissipated. If headache, dizziness in nausea develops during or after use, get out of contaminated area at once. Get medical attention.

WEAR clean clothing daily, wash all contamirated clothing with scap and hot water before reluse

Wash hands arms and face thoroughly with coap and water before eating or smoking.

DO NOT contaminate feed and foodstuffs. This part is paisonous to fish and wildlife. Keep at of lakes, pends and streams. Do not apply in any manner not specified on the label

Birds feeding on treated areas may be killed not apply where runoff is likely to occur lie not contaminate water by cleaning of equipment, or imposal of wastes.

Do not feed treated forage to dairy animals or animals being timished for slaughter.

Do not apply when weather conditions favor drift of spray or dust from areas reated

DESTROY all containers when empty by punch or holes in them and purying with wastes. Do to been

himse equipment and contenders and dispose towastes by purying to more pounds own from water supplies

NOT for use a stronge man of and the home

The The Table of the property of the control of the


#### ACTIVE INGREDIENTS:

Toxarhene	37.0°c
"C,O dimethyl O p nitrophenyl thiophosphate	37.0°
Xylene (Aromatic Petroleum Distillate) INEXT INGREDIENTS TOTAL	22.3° ; 3.7° c 100.0° ;

\*Technical chlorinated Camphene (Chlorine Content 67% to 69%)
\*Methyl Parathion

Each gallon of Cotton States Ketokil No. 44 contains 4 lbs. toxaphene plus 4 lbs. Methyl Parathion. Used as directed will control the following insects: Thrips, Fleahoppers, Plant bugs, Bollweevils, Bollworms, Cotton Leafworms, Aphids and Corn earworm.







#### ANTIDOTE FOR METHYL PARATHION

If SWALLOWED: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quite. Call a physician IMMEDIATELY.

IF ON SKIN. In case of contact remove contaminated clothing and immediately wash skin with soap and water.

#### PHYSICIAN'S NOTE

stimulation of the parasympathetic nervous system. Repeated use of a parasympathetic nervous system. Repeated use of a parasympathetic represent such as atropine to the limit of patient's tolerance, is indicated to relieve symptoms. The administration of oxygen is helpful. Never give morphine to patient's suspected of Parathion poisoning. 2 PAM is also antidetal and may be a tiministered in an in this with afficient.

Man itactured by



#### CWORKER TOMPTON

#### GENERAL DIRECTIONS

Prepare a spray mixture of Ketokil No. 44 a fillows:

- Add about half the total amount of water to be used to the spray tank
- Start agitator or pump bypass and add proper amount of Ketokil No. 44
- 3 Continue mixing and add remaining water

Directions for Use on COTTON

Ketokil No. 44 may be applied by either ground equipment or airroraft. For ground equipment, dilute in sufficient water to give adequate coverage. For aerial applications dilute in 3 to 4 gallons of water per acre.

Early Season Program: For general early season control, of plant bugs, thrips and flea hoppers, use .5 to 1 pint Ketokil No. 44 per acre. Make two or three applications at 7 d. intervals starting at the two-leaf stage.

Mid Season Insects Overwintering boll weevils, plant bugs, aphids, and cotton leaf worms are controlled with 1 to 2 pints Ketokil No. 44 per acre. Make repeat treatment at "liay intervals as innect counts indicate they are necessary.

Late Season Control. Use 2 to 4 pints per acre to control boll weevils bollworms and aphids. Use the higher rate for heavy boll worm outbreaks. Repeat applications at 5 to day intervals is insect counts indicate the necessary.