

CAUTION

Keep Out Of Reach Of Children

Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near food or food products. Keep children and domestic animals off treated areas until this material has been washed into the soil.

ANTIDOTE: If taken internally, induce vomiting and call physician at once.

Do not use or store near heat or open flame.

Use only with adequate ventilation.

Avoid contact with eyes and skin.

In case of contact, flush skin or eyes with plenty of water; for eyes get medical attention.

Wash thoroughly after handling.

Laundry clothing before re-use.

Avoid contamination of seed, feed and foodstuff.

In case of fire use water spray, foam, dry chemical or CO₂.

Flush spill area with water spray.

DIRECTIONS FOR USE

For Soybeans Only

Broadcast (40" Row Spacing)

Sandy Loam	Not Recommended
Silt Loam Soils	1/2 to 3/8 gallon
Heavy Clay Soils	2/3 to 3/4 gallon

12" Band

Silt Loam Soils	1-1/4 to 1-1/2 pints
Heavy Clay Soils	1-1/2 to 2 pints

1339-06934

LAY-LO

Lay-Lo is a liquid herbicide combination of Lasso® and LOROX® for pre-emergence weed control in soybeans.

ACTIVE INGREDIENTS:

*Alachlor	30.0%
**Linuron	15.0%
INERT INGREDIENTS	55.0%
	100.0%

*Contains 3 pounds 2-chloro-2', 6'-diethyl-N-(methoxymethyl) acetanilide per gallon.

**Contains 1.5 pounds [3-(3,4-dichlorophenyl)-1-methoxy-1-methylurea] per gallon. (equivalent to 3 lbs. LOROX per gal.)

Store above 32° F. to keep from freezing.

®LOROX is a registered trademark of E. I. DuPont de Nemours & Co.

®Lasso is a registered trademark of Monsanto Company.

Net Contents 1 U. S. Gallon
128 Fluid Ounces

Shake Well
Before Pouring



Agitate Well
While Spraying

USE LAY-LO ONLY ON SOYBEANS

BROADCAST (overall treatment) Apply the Recommended Rate of Lay-Lo in 20 or more gallons of water per acre on the soil surface after planting and before emergence of either crop or weeds. In areas of heavy grass or broadleaf weed infestations, use the higher rates in the recommended ranges.

DO NOT USE ON SAND OR LOAMY SAND AS CROP INJURY MAY OCCUR.

NOTICE: Seller warrants that the product conforms to its chemical description, and is reasonably fit for the purposes stated on the label when used in accordance with directions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

HOW TO MIX

Observe warning and handling precautions on label before mixing.

To achieve the best spray mix, proceed as follows:

1. Shake Lay-Lo mixture vigorously before using.
2. Fill tank one-half to three-fourths full with water.
3. Add the proper amount of Lay-Lo mixture to tank while agitating.
4. Fill the remaining portion of tank with water.

NOTE: Keep product in suspension at all times via vigorous agitation. If spray mixture is allowed to settle, thorough agitation is required to suspend the mixture before resuming spraying.

WEEDS CONTROLLED

Grasses	Broadleaf Weeds
Barnyardgrass	Giant Foxtail
Brachiaria	Yellow Foxtail
Crabgrass	Goosegrass
Fall Panicum	Red Rice
	Witchgrass
	Buttonweed (velvet leaf)
	Carpetweed
	Common Ragweed
	Smartweed
	Florida Fuzzley
	Teaweed
	Lambquarter
	Purslane
	Caroleweed (carelessweed)

WILL NOT CONTROL

Established Bermudagrass Establish'd Johnsongrass

NOTE: Do Not incorporate Lay-Lo as incorporation can cause crop injury.

Do Not use Lay-Lo in crops other than soybeans.

WHEN TO APPLY

Apply Lay-Lo mixture pre-emergence after planting and before emergence of either crop or weeds.

HOW TO APPLY

Lay-Lo tank mixture can be either broadcast or band applied. Best results are obtained when rainfall occurs within 3 to 7 days after application.