

DIRECTIONS

STAPLES — preplanting soil preparation: For the control of wire-worms, white grubs, Japanese beetle larvae, and white-fringed beetle larvae on potatoes, apply three quarts of **STAPLES** 8 lb. **CHLORDANE** per acre in sufficient water to insure coverage. This is the equivalent of 6 lbs. of technical Chlordane as called for in the 1963 Maine Potato Pest Control Bulletin. Apply **STAPLES CHLORDANE** prior to seeding and thoroughly work the soil to a depth of three or four inches immediately after application. Agitate water in sprayer while adding Chlordane. Destroy used containers to prevent re-use.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, that neither this warranty nor any other warranty of **MERCHANTIBILITY** or **FITNESS** for a **PARTICULAR PURPOSE**, express or implied, extends to the use of the product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

1345-3

STAPLES 8 LB CHLORDANE

ACTIVE INGREDIENTS:

Technical Chlordane*

Petroleum Distillate

INERT INGREDIENTS

TOTAL

*Equivalent to 42.4% Octachloro-4, 7-methanotetrahydroindane and 28.2% related compounds.

CONTAINS 8 LBS. TECHNICAL CHLORDANE PER GALLON

U.S.D.A. Registration No. 1345-3

WARNING

Keep Out of Reach of Children

Other precautions, first aid treatment, and antidote are on right side of label.

Net Contents 1 Gallon

Manufactured By

C. W. Staples, Inc.

PRESQUE ISLE, MAINE

STAPLES 8 LB. CHLORDANE

ACTIVE INGREDIENTS:

Technical Chlordane*	70.6%
Petroleum Distillate	22.4%
INERT INGREDIENTS	7.0%
TOTAL	100.0%

*Equivalent to 42.4% Octachloro-4, 7-methanotetrahydroindane and 28.2% related compounds.

CONTAINS 8 LBS. TECHNICAL CHLORDANE PER GALLON

U.S.D.A. Registration No. 1345-3

WARNING

Keep Out of Reach of Children

Other precautions, first aid treatment, and antidote are on right panel.

Net Contents 1 Gallon

Manufactured By

C. W. Staples, Inc.

PRESQUE ISLE, MAINE

ADDITIONAL

Do not get in eyes, on skin or in shoes. Do not take internally. Do not get on clothing. Keep away from heat and open flames. Do not apply to any water. Do not reuse empty container.

FIRST

EXTERNAL — In case of spillage on clothing, wash with water.

INTERNAL — Emetic of 1 table-spoon of 1% solution in water. Call a physician.

SYMPTOMS

Recognizable symptoms of acute intoxication are usually not as severe as those in general are similar and slowly progressive. Symptoms include loss of appetite, loss of weight, tremors, convulsions and eventual death.

NOTE TO ATTENDING PHYSICIAN

primary symptomatic. The following are the symptoms:

- 1- An attempt should be made to induce vomiting by use of sedatives. If vomiting does not occur, emetics are: phenobarbital, 100 mg. or more; hydrated, Use at levels that maintain blood levels which do not interfere with essential functions.
- 2- Definite measures should be taken to prevent further oral feeding if patient is comatose.
- 3- In cases of intoxication by ingestion, gastric lavage should be avoided.
- 4- Provide a quiet place as free from external irritation as possible.
- 5- Supportive treatment consistent with the patient's nervous system impairment.