


VEGETABLES



BEANS- Use 1 quart per acre in 30-50 gallons of water for control of Mexican bean beetle, leaf-hopper, aphid, red spider, armyworm, leaf roller and leaf miner. Apply at 7 to 10 day intervals as required. Do not apply within 15 days of harvest. The allowable dosage is 1½ quarts per acre (0.8 lbs parathion).

BEETS- Use 1 quart per acre in 30-50 gallons of water for the control of webworm and aphid. Do not apply within 15 days of harvest (21 days if tops are to be used for food or feed). The allowable dosage of this brand of parathion is 3 pints per acre (0.8 lbs parathion).

SUGAR BEETS- Use this brand of parathion at the rate of 1 quart per acre in 30-50 gallons of water for sugar beet webworm. Apply when insects first appear and repeat at 7 day intervals as necessary. Do not apply within 15 days of harvest (21 days if the tops are used for food or feed). The allowable dosage is 3 pints (0.8 lbs parathion) per acre.

CARROTS- Use ¾ pints per acre in 30-50 gallons of water for control of aphid. Do not use within 15 days of harvest. The allowable dosage is 2 quarts (1 lb parathion) per acre.

SQUASH AND MELONS- For the control of cucumber beetle, melon worm, serpentine leaf miner, aphid, and pickle-worm use 1 quart per acre in 30-50 gallons of water. Do not apply to squash within 15 days of harvest. Do not apply to melons within 7 days of harvest. The allowable dosage is 1 quart (0.5 lbs parathion) per acre.

PEAS- For the control of aphid, use 1 quart per acre in 30-50 gallons of water. Do not apply within 10 days of harvest. Where pea vines are used for forage, do not apply within 15 days of harvest. The allowable dosage is 1 quart per acre (0.5 lbs parathion).

POTATOES- Use 1 quart of this brand of parathion per acre in 30-50 gallons of water for control of flea beetles, leaf hopper and aphid. Do not apply within 5 days of harvest. The allowable dosage is 2 quarts per acre (1.0 lb of parathion).

TOMATOES- For tomatoes raised as field crops, use 1 quart per acre in 30-50 gallons of water for control of flea beetle, leafhopper and aphid. Do not apply within 10 days of harvest. Allowable dosage is 2 quarts (1.0 lb parathion) per acre.

CORN- For the control of European corn borer use 2 quarts per acre in 30-50 gallons of water. Apply first spray when 50% of eggs are in the blackhead stage. Make three applications 5 to 7 days apart. Do not apply within 12 days of harvest or cutting for forage. The allowable dosage is 2 quarts (1.0 lb parathion) per acre.

Original per acre

Left panel

DIRECTIONS FOR USE

----- COMPLETE COVERAGE SPRAY

Ground equipment recommendations call for 30-50 gallons of water per acre. Airplane recommendations call for five gallons of water per acre.

FRUITS

APPLES- Use 2 to 3 quarts Staples Parathion per acre in 30-50 gallons of water for the control of aphid, red-banded leaf roller, codling moth, scale insects and leaf miner starting at pre-pink stage and after petal fall. If infestation is not controlled, continue at 10-14 day intervals as needed. Do not apply to McIntosh type apples. The allowable dosage that may be applied is 3 gallons per acre (6.0 lbs parathion). Do not apply within 14 days of harvest.

BLUEBERRIES- For the control of maggots and thrips, use at the rate of 2 pints per acre in 30-50 gallons of water making applications before fruit sets or after harvest. The Allowable dosage is 2½ pints per acre (0.6 lbs parathion). Do not apply within 14 days of harvest.

PEACHES- Use 2 quarts per acre in 30-50 gallons of water for the control of plum curculio, oriental fruit moth, scale insects, cat-facing insects, green peach aphid and fruit tree leaf roller.


FOR AREAS OTHER THAN CALIFORNIA: Full coverage spray control of scale insects- Do not apply within 14 days of harvest. Do not apply more than 5 lbs actual parathion per acre (2½ gallons Staples Parathion). The allowable dosage is 2 gallons per acre (4.0 lbs parathion).

FOR CALIFORNIA: Do not apply within 21 days of harvest.

Do not apply more than once after bloom. Do not apply more than 2½ lbs actual parathion (5 quarts of this brand of parathion) per acre between January 1 and Harvest. The allowable dosage is 5 pints per acre (2.5 lbs parathion).

PLUMS AND PRUNES- Use 2 quarts per acre in 30-50 gallons of water for the control of plum curculio, spider mite, aphid, leaf roller, leafhopper and bud moth. Do not apply within 14 days of harvest. The allowable dosage is 2 gallons per acre (4 lbs parathion).

STRAWBERRIES- Use 2 quarts per acre in 30-50 gallons of water for the control of leaf roller and red spider mite, making applications at 7-10 day intervals as required. Do not apply within 14 days of harvest. The allowable dosage is 1½ quarts (0.8 lbs parathion) per acre



CABBAGE- Use 1 pint per acre in 30-50 gallons of water for the control of diamond-back moth, imported cabbage worm, army worm, aphid and thrip. Do not apply within 7 days of harvest in which case only the allowable dosage is 1 pint per acre (0.25 lbs parathion). Do not apply within 10 days of harvest in which case the allowable dosage is 1 quart per acre (0.5 lb parathion).

CUCUMBERS- For the control of cucumber beetle, serpentine leaf miner, aphid and pickle-worm, use 1 quart per acre in 30-50 gallons of water. Do not apply before plants begin to vine. Do not apply within 15 days of harvest. The allowable dosage is 1 quart (0.5 lbs parathion) per acre.

FORAGE CROPS- For the control of aphid, grasshoppers, army worm found on wheat, oats, barley, clover, grass, hay and alfalfa use 1 quart per acre in 30-50 gallons of water. Do not apply within 15 days of cutting. The allowable dosage is $1\frac{1}{2}$ quarts per acre (0.8 lbs parathion) for alfalfa, grass and hay. For wheat, oats, barley and clover, the allowable dosage is 3 quarts (1.5 lbs parathion) per acre. **NOTE ON SPRAYING PASTURES WITH PARATHION.** Do not apply within 15 days of pasturing.

ACCEP

MARCH 26-1968

1395-8
IS ATTACHED COMMENT

STAPLES 25% PARATHION

(Back Panel)

WARNING

Poisonous if swallowed, inhaled or absorbed through the skin. Do not get in eyes, on skin or on clothing. Keep away from heat or open flame. Wear a mask or respirator of a type approved by the U. S. Department of Agriculture for parathion protection.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label. Do not store or use in or around the home.

Do not apply when weather conditions favor drift from areas being treated. Do not contaminate water by cleaning equipment or disposal of wastes. Do not reuse empty containers. Destroy containers when empty.

Wash hands and face thoroughly with soap and water before eating or smoking. Wear protective clothing, rubber gloves and goggles. Keep all unprotected persons out of treated area or vicinity where there may be danger of drift. Treated areas should not be reentered until volatile residues have dissipated.

ANTIDOTE STATEMENT

SYMPTOMS OF POISONING- A sense of "tightness" in chest; stomach pains; vomiting and diarrhea.

ANTIDOTE- Internal: Give a teaspoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have patient lie down and keep quiet. Call a physician at once.

ANTIDOTE- External: If the skin is accidentally contaminated, wash immediately with soap and warm water.

TO PHYSICIAN- Compound exerts muscarine-like effects, that is, stimulation of the parasympathetic nervous system. A parasympathetic depressant, such as atropine, is indicated to relieve symptoms. The administration of oxygen is helpful.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

ACCEPTED

MAR 26 1968

1345-8