

RECOMMENDED USES

LIVESTOCK—

RANGE AND FEEDER CATTLE: For control of Sarcoptic Mange, Chorionic Mange, Psoroptic Scab, Cattle Lice, Ear Tick, Lone Star Tick, Winter Tick, Cattle Fever Tick and Horn Flies—Use 1 gallon of this product to 150 gallons of water and stir or agitate well to form a creamy emulsion. Spray or dip animals thoroughly to assure penetration to the skin. For spraying operation it is advisable to use a high pressure sprayer at a minimum of 250 pounds pressure with constant agitation of spray mixture in the tank. If spray-dip machines are used we recommend that the manufacturers recommendations and directions for operation be used. It is essential that suction strainers be removed and cleaned as needed to remove hair and debris to permit recycling of the spray mixture.

For use in dipping vats the same concentration is used as given above. Before dipping operation thoroughly mix the vat contents by using a pole or bucket attached to a rope. Addition of a wetting agent is advisable if the water is known to be extremely hard and highly alkaline. Standard methods used for determining dipping vat concentration of toxaphene emulsions may be used for this product. **NOTE:** Do not dip calves under three months of age. Avoid treating sick or overheated animals. Avoid spray or dip treatment during stormy or cold weather. If treating in warm weather provide shade for animals after treatment. Water animals before dipping or spraying to prevent them from drinking the dip or puddles from the spray. In case of quarantine where calves require treatment spray very lightly. Do not treat animals within 28 days of slaughter.

BACK RUBBERS: BEEF CATTLE—As an aid for the control of Lice and Hornflies, mix 1 gallon to 15 gallons of fuel oil or mixture of mineral oil and soak sack or cloth of burlap rubbing device or fill compartment of back rubber as required. Recharge back rubber as necessary. Mount or place back rubber where it is readily accessible to animals as they move to feed, water or pass through gate from one pen to another. Do not use on calves under six months of age or around dairy animals. Do not use as a spray directly on animals. Animals should not be permitted access to back rubbers within 28 days of slaughter.

ACCEPTED
MAY-28-1971
SELCO
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC CONTROL OF PESTS
REGISTERED UNDER PATENT NO. 1348-38
TO THE U.S. GOVERNMENT

Toxaphene 6 E Emulsifiable

For Control of Livestock Pests and Field Crop Insects

ACTIVE INGREDIENT:

Toxaphene*	59.00%
Aromatic Petroleum Derivative Solvent	31.33%
INERT INGREDIENTS	9.67%
TOTAL	100%

* Technical Chlorinated Camphene (Chlorine Content 67 to 69%)
CONTAINS 6-LBS. TOXAPHENE PER GALLON

WARNING KEEP OUT OF REACH OF CHILDREN!
See side panel for further Warnings!

WARNING

May be fatal if swallowed! Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Do not apply in dairy barns or on dairy animals. Do not use on beef cattle, sheep or goats within 28 days of slaughter. Do not use in poultry houses.

ANTIDOTE

INTERNAL: Give sodium chloride, 15 grams (1 tablespoon of table salt) in a glass of warm water and repeat until vomit is clear. **CALL A PHYSICIAN IMMEDIATELY.**

EXTERNAL: Wash with soap and water.

SHEEP
at the
animal
Lice—
150 gal
treat s
or goat

SMALL
Apply
ground
in a n
in spr
grazer
being
same
water.
first se
Do not
or anim

WARN
on trea
Do not
of was

NOTICE
burying
should
wastes

NOTICE
product

REGISTERED

4-28-1971
SELCO

THE FEDERAL INSECT AND
DISEASE CONTROL ADMINISTRATION
WASHINGTON, D. C. 20540
1348-38

Toxaphene 6 E Emulsifiable

Control of Livestock Pests and Field Crop Insects

ACTIVE INGREDIENT:

Toxaphene*	59.00%
Aromatic Petroleum Derivative Solvent	31.33%
INERT INGREDIENTS	9.67%
TOTAL	100%

Technical Chlorinated Camphene (Chlorine Content 67 to 69%)
CONTAINS 6-LBS. TOXAPHENE PER GALLON

WARNING KEEP OUT OF REACH OF CHILDREN!
See side panel for further Warnings!

WARNING
Is fatal if swallowed! Do not breath spray mist. Do not get on skin or on clothing. Wash thoroughly after using. Contamination of feed and foodstuffs. Do not apply in dairy or on dairy animals. Do not use on beef cattle, sheep or goats 28 days of slaughter. Do not use in poultry houses.

ANTIDOTE
INTERNAL: Give sodium chloride, 15 grams (1 tablespoon ble salt) in a glass of warm water and repeat until is clear. **CALL A PHYSICIAN IMMEDIATELY.**
EXTERNAL: Wash with soap and water.

RECOMMENDED USES

SHEEP AND GOATS: For control of Psoroptic Sheep Scab use at the rate of 1 gallon to 150 gallons of water. Thoroughly wet animals. For control of Sheep Ticks (Keds), Sheep Lice and Lice—Use 1 gallon to 300 gallons of water for dip and 1 gallon 150 gallons of water for a spray. Wet animals thoroughly. Do not treat sick animals. Do not treat milk goats. Do not use on sheep or goats within 28 days of slaughter.

SMALL GRAINS (Barley, Oats and Wheat): Army Cutworms — Apply at the rate of 1 qt. to 50 gallons of water per acre by ground spray equipment. For airplane application apply 1 qt. in a minimum of 5 gallons of water per acre. Make Application in spring when larvae are small and first seen. Do not feed or graze forage from treated grains to dairy animals or animals being finished for slaughter. For Grasshoppers: Apply at the same rate as for Army Cutworms using the same dilution of water. Make application in spring when nymphs are small and first seen, in hatching areas. Apply only one application per season. Do not graze or feed forage from treated grains to dairy animals or animals being finished for slaughter.

WARNING: This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on the label.

NOTICE: Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

NOTICE: Buyer assumes all risk of use, storage or handling of this product when not in strict accordance with directions given herewith.

USDA REG. NO. 1348-38

NET CONTENTS 1 QUART

Formulated By
SELCO SUPPLY COMPANY
EATON, COLORADO