

RECOMMENDED USES

CONTROL OF ELM BARK BEETLES: This product is recommended for spraying elm trees for control of elm bark beetles, the principal carriers of the fungus which causes Dutch Elm disease.

For Hydraulic Sprayers: Mix 8 gallons of product to 100 gallons of water. Apply sufficient spray to thoroughly wet all bark on trunk, limbs and twigs; 20 to 30 gals. of spray are usually required for a 50-foot elm tree.

For Mist Sprayers: Mix 5 gallons of product to 10 gallons of water. Thorough coverage of all bark surfaces is important; usually 2 to 3 gals. of spray are required to adequately cover a 50-foot elm tree.

Make application as a dormant spray treatment anytime after the elm trees lose their leaves in the fall and before the new leaves or flowers appear in the spring. Apply when temperature is above 40 degrees F. and there is no danger of freezing before the spray dries.

For further uses of this product consult your local State Agricultural Extension Service or State Experiment Station for additional information consistent with label directions and restrictions.

NET CONTENTS 1 GALLON

SELECO

ACCEPTED
MAY 27 1971
UNDER THE
FEDERAL
INSECTICIDE ACT

METHOXYCHLOR
24% EMULSIFIABLE

ACTIVE INGREDIENTS:

Methoxychlor, Technical	24%
Methylated Naphthalenes	60%
INERT INGREDIENTS	16%
Total	100%

24% methoxychlor, technical is equivalent to 1,1,1-tris-(2-methoxyphenyl)-1,1,1-trichloroethane and its other isomers and reaction products.
1 gallon contains 2 pounds of actual Methoxychlor.

CAUTION! Keep out of reach of Children.

Hazard if swallowed.
Avoid prolonged or repeated contact with skin. Avoid prolonged or repeated breathing of spray mist. Avoid contamination of food and feed. In case of contact, wash with soap and water.

LIVESTOCK: (Beef Cattle) — For Hairy Control — Use 1 quart of product to 12½ gals. water. Apply 2 quarts of spray per mature animal and proportionally less for smaller ones. Note: Use as a spray only. Do not use in dipping vats. Do not spray Dairy Animals.

GRAIN STORAGE BINS: To clean up infestations in empty storage bins, use 1 gallon of product to 15 gallons of water applied at the rate of 2 gallons of spray per 100 square feet before products are stored. This aids in control of Cabinet Flour Grain Beetle, Saw-toothed Grain Beetle, Lesser Grain Borer, Red Flour Beetle, Long-headed Flour Beetle, Rice Weevil, Gummy Weevil, Confused Flour Beetle, Foreign Grain Beetle, and Hairy Fungus Beetle. Note: Do not add grain to bin until at least 24 hours or until walls have dried out thoroughly.

If exposed to prolonged cold, place in warm storage (50 to 60 degrees F.) for several hours and agitate before using.

DO NOT use, pour, spill or store near heat or open flame.

NOTICE: Containers should be disposed of by punching holes in them and burying with wastes in non-crop lands away from water supplies.

NOTE: Toxic to fish. Keep out of lakes, ponds and streams.
NOTICE: Higher temperatures will risk of a reaction occurring if the product when not in strict accordance with directions.

U.S.D.A. REG. NO. 1348-48

FORMULATED BY
SELECO SUPPLY COMPANY
EASTON, COLORADO