

ACCEPTED

APR 1972

UNDER THE
FUNCTION
FOR LABELING OF
ED UNIFORM

1348-72

SELCO MALATHION 5% DUST MN-5

EPA Reg. No. 1348-72

ACTIVE INGREDIENT:
Malathion^o 5.00%
INERT INGREDIENT 95.00%

*0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate.
RECOMMENDED FOR CERTAIN INSECTS ON VEGETABLES,
ORNAMENTALS AND LIVESTOCK AND PETS

DIRECTIONS FOR USE ON VEGETABLES AND ORNAMENTALS

- BEANS:** For control of Mexican bean beetle apply at rate of 25 to 30 lbs. per acre. Repeat application as needed but not within 1 day before harvest.
- CABBAGE:** For control of cabbage worm, cabbage looper and cabbage aphid apply at the rate of 30 lbs. per acre. Repeat application as needed but no later than 7 days before harvest.
- CUCUMBERS:** For control of melon aphids and thrips apply dust at rate of 20 to 25 lbs. per acre. Repeat application as needed but do not apply within 7 days of harvest. Do not apply dust to cucurbits unless plants are dry.
- ORNAMENTAL PLANTS:** For control of aphids and spider mites dust plants thoroughly. Repeat treatment as necessary.

DIRECTIONS FOR USE ON LIVESTOCK AND PETS

- BEEF CATTLE (Only):** For control of lice make a thorough application of dust to animals. Repeat application in 10 days and thereafter as needed. Do Not Treat Calves Under One Month of Age. For the control of hornflies on beef and dairy cattle apply 3 table-spoons of 5% dust per animal. Apply the dust on the back and neck of the animals and repeat application at 10-14 day intervals. Do not apply to dairy cattle later than five hours before milking or during milking. Do not use sprays or dip dairy animals in malathion. Do not contaminate water troughs.
- POULTRY:** For control of northern fowl mite and poultry lice dust individual birds applied by means of a shaker can or with a rotary hand duster. Repeat in 4 to 8 weeks or when necessary. Avoid contaminating feed, feeding troughs and water fountains with dust.
- POULTRY HOUSES:** For control of northern fowl mite, chicken red mite and poultry lice, apply 1 lb. of dust per 50 to 60 sq. ft. of litter and floor space and to nests, roosts and adjacent areas. Dust may be applied with a rotary hand duster or sprinkling from a can. Repeat as necessary. Avoid contaminating feed, feeding troughs, and water fountains with dust.
- DOGS & CATS:** For control of fleas on dogs and cats apply dust using enough to treat the animals thoroughly. Apply with a puff duster or with a shaker-type dispenser. If animal has free run, treatments may be made every second or third week. Avoid unnecessary treatment by close examination for fleas. If animal is confined to an area free of fleas, one treatment may be sufficient. DO NOT treat kittens or puppies under 4 weeks of age.
- ANIMAL QUARTERS:** For control of fleas in dog and cat kennels or sleeping quarters apply dust at rate of 1 to 2 lbs. per 1000 sq. ft. using a hand duster. Remove manure or debris before treatment.

CAUTION! *Keep out of reach of children!*

CAUTION: Avoid breathing of dust. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs.

This product is highly toxic to bees exposed to direct treatment. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product will kill fish. Keep out of any body of water.

Do not re-use container. Destroy when empty.

NOTICE: Buyer assumes all risk of use, storage or handling of this product when not in strict accordance with directions given herewith.

Formulated
By— **SELCO SUPPLY COMPANY** 4 Pounds
Net Weight
EATON, COLO. 80615