

1348-84

DIRECTIONS

GRAIN ELEVATORS, FEED STORAGE, FARM BUILDINGS, FOOD PROCESSING PLANTS, AND OTHER BUILDINGS
 Flea Beetles, Grain Weevils, Rice Weevils, Grain Beetles, Carpet Beetles, Grain Mites, etc. Dilute 1 to 10. Before attempting to place product, thoroughly clean and vacuum or all least vacuum corners and crevices, remove all debris, dirt, grain, and other infested food products from the area to be treated.
 Insecticide: Spraying of area with water and detergent or other water based detergent on walls, on cracks and partitions, on floors under equipment, and on other places where insects are likely to be present. Apply insecticide spray at the rate of 1 to 2 gallons per 100 sq. ft. of wall area. Use portable equipment in areas. Repeat cleaning operations and spraying of walls as needed. Do not use vacuum suction which could be injurious to walls.
 Insects: Treat entire area with insecticide every 30 days if grain is present, or every 60 days if no grain is present. Do not use in buildings where grain is stored. For use with the Animal Quarantine and Control Laboratory, the contents of this container should be used in treated areas. Ventilate the area after treatment.
 Cockroaches, Waterbugs, Ants, Silverfish, Cheese Mites, and Certain Spiders: Dilute 1 to 2 with water and apply diluted emulsion at the rate of 1 gallon per 75 square feet of surface to be sprayed. Spray thoroughly into all hiding places, cracks, crevices, and under and behind equipment, shelves, desks and counters. Repeat treatments at regular intervals or as often as necessary to keep insects under control. **CAUTION:** Protect pets, birds, and other farm animals before treatment.

NET CONTENTS 1 GALLON

• PYRENONE

EMULSIFIABLE CONCENTRATE

EPA Reg. No. 1348-84

ACTIVE INGREDIENTS	
Technical Pyrenone Emulsion**	10.00%
Paraffin	1.00%
Petroleum Solvents	2.00%
INERT INGREDIENTS	
TOTAL	13.00%

**Equivalent to 50% of total weight of emulsion (petroleum solvents and other inert ingredients)

*Registered trademark of U.S. Industrial Chemicals, Inc.

CAUTION: HARMFUL IF SWALLOWED!
Keep Out of Reach of Children

Livestock Uses: Use on Horses, Cattle, Goats, Hogs and Sheep. Dilute 1 to 10. Apply liberally on legs, neck, and body. Use at a rate of 1 to 4 oz. for sheep, goats, and hogs. Dilute 1 to 10 for Sheep, Goats. Dilute 1 to 10 with water. Apply as a spray or by drench to wet all portions of body. Repeat as needed.

Use on Animals: Use on Pasture Horses, Pasture Cattle, etc. Dilute 1 to 10. Apply liberally on legs, neck, and body. Use at a rate of 1 to 4 oz. for sheep, goats, and hogs. Dilute 1 to 10 for Sheep, Goats. Dilute 1 to 10 with water. Apply as a spray or by drench to wet all portions of body. Repeat as needed. For feeding control dilute 1 to 10 with water.

CATTLE - HORSES: Flea Fleas, Stable Fleas and Lice. Dilute 1 to 10 with water. Apply as a spray at the rate of 1 to 4 oz. per animal. For control of lice, use 1 to 4 oz. per animal. Dilute 1 to 10 with water. Apply to the legs and under the body. Repeat treatments as needed. For control should be made monthly every 30 days.

CATTLE - HORSES: Flea Fleas and Mites. Dilute 1 to 10 with water. Apply as a spray at the rate of 1 to 4 oz. per animal. For control of lice, use 1 to 4 oz. per animal. Dilute 1 to 10 with water. Apply to the legs and under the body. Repeat treatments as needed. For control should be made monthly every 30 days.

NOTICE: Pyrenone is a registered trademark of U.S. Industrial Chemicals, Inc.

Do not use in container in which other agents are present.

Formulated By
ISCO SUPPLY COMPANY
BATON, LOUISIANA 70801