

1348-93

ACCEPTED
U.S. DEPARTMENT OF AGRICULTURE
FEDERAL BUREAU OF INVESTIGATION
REGISTERED UNDER 1348-93 SUBJECT TO FEDERAL CONTROL

NOTICE: Rinse equipment and containers and dispose of wastes by burning in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying them with wastes.

NOTICE

Buyer assumes all risk of use, storage or handling of this product when not in strict accordance with directions given herewith.

SELCO

LINDANE 25% WETTABLE

ACTIVE INGREDIENTS:	
Gamma Isomer of Benzene Hexachloride (from Lindane)	25.00%
INERT INGREDIENTS	75.00%
TOTAL	100.00%

USDA REG. NO. 1348-93

WARNING! Keep out of reach of children!
See side panel for additional warnings!

MAY BE FATAL IF SWALLOWED! MAY BE ABSORBED THROUGH THE SKIN! MAY CAUSE IRRITATION OF SKIN AND EYES! Do not breathe dust or spray mist. Avoid contact with skin and eyes. Avoid contamination of food and foodstuffs. Wash thoroughly after handling and before eating or smoking. In case of skin contact, wash with plenty of soap and water. For eyes, flush with plenty of water for at least 15 minutes and get prompt medical attention. Do not use treated seed for animal feed or food processing. When treating seed, wear respirator to prevent inhaling dust. Wear gloves and tight fitting clothing to prevent skin contact. Use clean clothing daily.

Toxic to Fish and Wildlife. Keep out of lakes, ponds and streams!

NET CONTENTS 1 POUND

LIVESTOCK — GENERAL DIRECTIONS and CAUTIONS

DO NOT use this material on emaciated or lactating animals. Animals should have shade following dipping or spraying in warm weather. DO NOT treat animals in cold stormy weather. DO NOT treat sick animals. DO NOT apply to dairy animals. DO NOT treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter. Do not treat animals under 3 months of age. Spray or dip applications to young animals (over 3 months of age) should be limited to the 0.03% Lindane concentration. DO NOT dip animals within 60 days of slaughter or spray animals within 30 days of slaughter.

CATTLE (Beef): Lice and Ticks—Use 1 lb. in 100 gal. of water. Use 1 gallon of spray per adult animal for thorough coverage, and repeat in 2 to 3 weeks if necessary. For Sarcoptic Mange on Beef Cattle use 1½ lbs. in 100 gal. water, applying 2 gals. of spray to adult animal. A second application 2 to 3 weeks later is required for a complete cleanup. Do not use on calves under 3 months of age.

PIGS: Lice and Sarcoptic Mange—Use 1 to 2 lbs. in 100 gal. water. Wet animal thoroughly. Apply 2 to 4 qts. per adult animal. Repeat in 2 to 3 weeks if necessary.

FORMULATED BY
SELCO SUPPLY COMPANY
EATON, COLORADO