



This product is especially formulated to be diluted with water to be used as a livestock spray. It can also be diluted with distillate or diesel fuel and used in Livestock Oilers and Back Rubber devices.

BEEF CATTLE: For the control of Lice, Ticks and Sarcoptic Mange Mites on mature cattle, use at the rate of 1 quart to 37 gallons of water. Spray animals thoroughly applying at least 2 gallons of finished spray per head to larger animals for complete coverage. Repeat treatment in 10 to 14 days if necessary. Avoid treating animals in cold, stormy weather. Avoid treating overheated animals.

For the control of Horn Flies and Mosquitoes, use at the rate of 1 quart to 37 gallons of water applying thorough coverage to top line and side portions of animals body. Treatment for the above summer pests require at least 2 quarts of finished spray per animal for each treatment at 2 to 3 week intervals. Do not apply to cattle within 30 days of slaughter. Avoid treating animals in cold, stormy weather. Avoid treating overheated animals.

SHEEP, GOATS and HOGS: For the control of Lice and Ticks on Sheep and Goats and Sarcoptic Mange Mites and Lice on Hogs, use at the rate of 1 quart to 50 gallons of water. Spray animals thoroughly. Repeat treatment in 10 to 14 days if necessary. Do not apply to sheep, goats and hogs within 30 days of slaughter. Avoid treating animals in cold, stormy weather. Avoid treating overheated animals. Do not use on dairy goats. Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter.

FOR USE IN A LIVESTOCK OILER OR BACK RUBBER DEVICE—Control of Horn Flies: Dilute SELCOTOX at the rate of 1 gallon to 10 gallons of distillate or diesel fuel. Place diluted mixture in oil reservoir or on burlap or oiler device and keep wet at all times. Place oiler equipment where animals have free access to rubbing and near watering facilities or cattle crossings wherever cattle congregate. Do not use for dairy cows or calves under 6 months of age. Beef cattle should not be exposed to the backrubber treatment within 30 days of slaughter. Avoid treating animals in cold, stormy weather. Avoid treating overheated animals.

EATON & SEVERANCE, COLORADO



SELCOTOX LIVESTOCK SPRAY

TOXAPHENE AND LINDANE

| | |
|--|---------|
| ACTIVE INGREDIENTS: | |
| Toxaphene (Technical Chlorinated Camphene containing 67 to 69% Chlorine) | 42.63% |
| Gamma Isomer of Benzene Hexachloride from Lindane | 1.80% |
| Aromatic Petroleum Derivative Solvent | 45.71% |
| INERT INGREDIENTS | 9.86% |
| TOTAL | 100.00% |

Contains 4 lbs. Toxaphene and .18 lbs. Lindane per gallon

EPA REG. NO. 1348-108

EPA EST. NO. 1348 — CO — 1

WARNING: Keep out of reach of children!

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

See Side Panel for Additional Warnings.

WARNING

MAY BE FATAL IF SWALLOWED—May cause irritation of skin or eyes. May be absorbed through the skin. Avoid skin contact. Wash skin thoroughly with soap and water immediately after contact or use. Keep clothing free from residues. Remove and wash contaminated clothing. Avoid inhaling spray mist or getting it in the eyes. For eyes get medical attention. Flush with water for 15 minutes. Use respirator and goggles if necessary for protection. Combustible Mixture. Don't keep poisons where food is handled or stored. Avoid food contamination.

WARNING: This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. Do not use on Dairy Animals or in Farm Buildings. Do not use on animals under 6 months of age. Do not treat sick animals or spray animals within 30 days of slaughter. Do not use this product on dogs, cats, or other household pets nor on dairy animals or poultry. Do not use on newborn (very young) animals or calves under 6 months old. Do not use on hogs within 2 weeks of farrowing nor within 3 weeks after farrowing. Do not use in dairy barns, milk rooms, or poultry houses.

NOTICE

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning. (Keep out of Smoke).

NOTICE

Buyer assumes all risk of use, storage or handling of this product when not in strict accordance with directions given herewith.

NET CONTENTS 1 GALLON

FORMULATED BY

SELCO SUPPLY COMPANY