

DIRECTIONS FOR USE

FLOWERS: Roses, asters, carnations, chrysanthemums, marigolds, zinnias, azaleas, gladioli, snapdragons, peonies, petunias, phlox, delphinium—TO CONTROL—Leafhoppers, aphids, fleahoppers, rose chafer, lace bug, flea beetles, rose slugs (sawflies), and tent caterpillar, apply at the rate of 2 Tbl. to 1 gallon of water (1 pt. to 16 gals. of water). Make first application when insects first appear and repeat application as necessary.

ORNAMENTALS, EVERGREENS and SHADE TREES: TO CONTROL—Aphids, red spider mites, cankerworms and other insects listed for flowers, use 2 Tbl. to 1 gallon of water (1 pt. to 16 gallons of water).

For scale insects double the above dosage. Make first application when insects first appear and repeat treatment as necessary. NOTE: May cause injury to ferns, maidenhair, pteris and crassula.

VEGETABLES: Beans — Mexican bean beetle, Bean leaf beetle and Flea beetle. Potatoes — Colorado potato beetle, flea beetles and leafhoppers. Tomatoes — Leafhoppers, tomato fruitworm and red spider. Cabbage — Cabbage worms. Peas — Aphids. Peppers — Aphids. Apply this product at the rate of 2 Tbl. to 1 gal. water (1 pt. to 16 gals. of water). On an acre basis use 1 gal. of this product to 128 gals. of water. Apply when insects first appear and repeat treatment as necessary. (Do not apply within 7 days of harvest).

EPA REG. NO. 1348-153



Garden & Floral Spray

KILLS MANY CHEWING AND SUCKING INSECTS

ACTIVE INGREDIENTS	25.00%
*Methoxychlor, Technical	12.50%
**Malathion	12.50%
INERT INGREDIENTS	75.00%
*12.50% methoxychlor, technical is equivalent to 10.94% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 1.56% other isomers and reaction products.	
**O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.	

CAUTION: KEEP OUT OF REACH OF CHILDREN!

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

HARMFUL IF SWALLOWED—Avoid breathing of spray mist. Avoid contact with skin; wash thoroughly after handling. Store away from children and domestic animals. Avoid contamination of feed and foodstuffs. THIS PRODUCT is highly toxic to bees exposed to direct treatment or residues on crops. Do not make treatment when bees are actively visiting the area.

NET CONTENTS 1 QUART

Agricultural Chemicals



Garden & Floral Spray

KILLS MANY CHEWING AND SUCKING INSECTS

ACTIVE INGREDIENTS	-----	25.00%
*Methoxychlor, Technical	-----	12.50%
**Malathion	-----	12.50%
INERT INGREDIENTS	-----	75.00%
*12.50% methoxychlor, technical is equivalent to 10.94% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 1.56% other isomers and reaction products.		
**O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.		

CAUTION: KEEP OUT OF REACH OF CHILDREN!

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

HARMFUL IF SWALLOWED—Avoid breathing of spray mist. Avoid contact with skin; wash thoroughly after handling. Store away from children and domestic animals. Avoid contamination of feed and foodstuffs. **THIS PRODUCT** is highly toxic to bees exposed to direct treatment or residues on crops. Do not make treatment when bees are actively visiting the area.

NET CONTENTS 1 QUART

Agricultural Chemicals

FRUITS: Apples, cherries, pears and plums—
TO CONTROL—Aphids, red spider, codling moth, plum curculio, leafhoppers, pear slug, apple maggot, cherry fruitworm, cherry fruit fly; use at the rate of 2 Tbl. to 1 gal. of water (1 pt. to 16 gals. of water). Apply when insects first appear (for codling moth begin spraying at petal fall stage) and repeat treatment at 10 day to 2 week intervals as necessary to maintain adequate control. Spray foliage on both bottom and top, also spray on fruit to maintain thorough coverage of insecticide. Do not apply within 7 days of harvest.

BERRIES: Blackberries, grapes, raspberries, gooseberries and strawberries — TO CONTROL — Aphids, leafhoppers, grape berry moth, grape leaf skeletonizer, red spider, and strawberry weevil, use at the rate of 2 Tbl. to 1 gal. of water (1 pt. to 16 gals. of water—on an acre basis use 1 gal. of this product to 128 gallons of water). Apply when insects first appear and repeat treatment as necessary. Do not apply within 21 days of harvest.

This product is Toxic to Fish. Keep out of Lakes, Ponds and Streams.

Do not reuse container. Destroy when empty. Apply this product only as specified on this label.

DO NOT use, pour, spill, or store near heat or open flame.

NOTICE: Buyer assumes all risk of use, storage or handling of this product when not in strict accordance with directions given herewith.

FORMULATED BY