

21

SILCO

CAPTAN 7.5% DUST

(AN AGRICULTURAL FUNGICIDE)

ACTIVE INGREDIENT.	
Captan (N-trichloromethylthio-4-cylohexen-1,2-di carboximide)	7.5%
INERT INGREDIENTS	92.5%
Total	100.0%

U. S. Reg. Pat. Nos. 2,552,770, 2,552,771, 2,552,776

EPA REG. NO. 134867

EPA EST. NO. 1348-CO-1, 2

RECOMMENDATIONS

BEANS, PEAS: Soil treatment—Damp-off (beans only), Root Rot—Work into top 3 or 4 inches of soil before planting. Apply 50 to 60 lbs. per acre.

CELERY: (Plant Bed Only)—Damp-off, Pink rot—Apply 40 to 50 lbs. per acre when all seeds have germinated. Repeat if necessary. Cucurbits (cantaloupe, cucumber, melon, squash): Angular Leaf Spot, Anthracnose, Downy Mildew—Start application when runners begin to form or at the first sign of disease. Apply at rate of 20 to 30 lbs. per acre on young plants increasing to 30 to 40 lbs. per acre on mature plants. Repeat at 7 day intervals.

POTATO SEED BENCH TREATMENT: Damp-off, Seed rot—Apply at rate of 2 lbs. per 100 lbs. pieces. Coat each potato seed piece evenly with a light dust. Treat pieces within 6 hours after cutting. If planting is delayed store treated seed pieces in open crates 2 or more days before bagging.

DO NOT use treated seed pieces for food or feed.

SOIL AND GREENHOUSE BENCH TREATMENT: Apply 2½ lbs. per 1000 sq. ft. (100 lb. per acre). Work into upper 3 or 4 inches of soil before planting. For damp-off and root rot diseases of seedlings and transplants of roses, other shrubs, flowers, lawn seed beds, and vegetables such as beans, peas, celery, cabbage, eggplant, pepper and tomato.

NOTE: In order that residues on food and forage crops will not exceed established tolerances use only at recommended rates, and do not apply closer to harvest than specified. Observe use limitations given on label for specified crops.

CAUTION:

Keep out of reach of children!

Avoid inhaling dust. Avoid contact with skin. Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

NOTICE: Containers should be disposed by punching holes in them and burying with wastes.

NOTICE: Buyer assumes all risk of use, storage or handling of this product, when not in strict accordance with directions given herewith.

NET WEIGHT 50 POUNDS

Formulated By **Selco Supply Co.** Eaton, Colorado

FEB 1975

Under the provisions of the Federal Insecticide, Fungicide, and Rodenticide Act