

**ACCEPTED**

JAN 31 1973

UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 ON ECONOMIC POISON REGISTERED  
 UNDER NO. 1348-173 SUBJECT  
 TO ATTACHED COMMENTS.

1348-173

SELCO

# SEVIN® 50% WP

## RECOMMENDED USES FOR INSECT PESTS OF FRUIT, VEGETABLES AND ORNAMENTALS

ACTIVE INGREDIENT: Carbaryl (1-naphthyl N-methylcarbamate) 50%  
 INERT INGREDIENTS .....50%  
 Total .....100%  
 U. S. Pat. No. 2,903,478

### CAUTION:

### Keep Out of Reach of Children!

**HARMFUL IF SWALLOWED OR INHALED** Do not take internally. Avoid breathing dust or spray mist. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed, feeding troughs, watering receptacles and milking utensils.

#### CAUTION IN USE

Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following spraying. Does not control spider mites but is compatible with all common miticides (except Bordeaux or Lime Sulfur mix).

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

Do not reuse container. Destroy when empty.

**NOTICE:** Buyer Assumes all risk of use, storage or handling of this product when not in strict accordance with directions given herewith.

EPA REG. NO. 1348-173

"Sevin" Trademark of the Union Carbide Corporation

**NET CONTENTS 8 OZS.**

**DIRECTIONS FOR USE**  
**VEGETABLE INSECT CONTROL:** For Home Gardens—Use 3 level tablespoonfuls to 1 gal. water. Keep spray mixture well agitated. Apply when insects first appear and repeat treatment as necessary. Spray thoroughly bottom and tops of plant foliage for optimum insect control.

**BEANS:** Mexican bean beetle, leafhoppers, flea beetles, western bean cutworm and stink bugs.

**CORN (Sweet):** Corn Earworm and Corn root worm adults—For larvae in whorl and foliage feeders apply to entire plant. Repeat as necessary. For protection of insects attacking silks and ears, apply at 3 to 8 day intervals starting when first silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage is necessary.

**MELONS, PUMPKINS, SQUASH, CUCUMBERS:** Pickleworm, melonworm, leafhoppers, flea beetles and squash bugs—Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in presence of very high humidity.

**POTATO, TOMATO, EGGPLANT AND PEPPER:** Colorado potato beetle, fleabeetles, leafhoppers, tomato fruit worm and tomato hornworm and grasshoppers.

DO NOT apply within 1 day of harvest on the vegetables listed above.

**CABBAGE, BROCCOLI, BRUSSELS SPROUTS CAULIFLOWER:** Corn earworm, cabbage worm and small cabbage loopers. For small cabbage loopers use in 5 to 7 day schedule. Do not apply within 8 days of harvest.

**ORNAMENTAL INSECT CONTROL:** Recommended concentrations can be safely used on a wide range of ornamental plants, flowers, shrubs and shade trees including roses, peonies, juniper, arbor vitae, elm, maple, oak, dogwood, birch and alder. Do not use on Boston Ivy. Use 2 lbs. to 1 gal. or 2 lbs. per 100 gallons of water when insects first appear. Repeat as needed for control of Bagworms, Box Elder Bug, Elm Leaf Beetle, Periodical Cicade, Flea Beetles, Lace Bugs, Leafhoppers, Leaf Rollers, Plant Bugs, Rose Aphid, Rose-Hug, Scale insects, Tent Caterpillars, Thrips (exposed), and Willow Leaf Beetles. For optimum control of scale insects application should be made in the crawler stage.

**LAWN INSECT CONTROL:** Ants, chinch bugs, earwigs, fall armyworms, fleas, leafhoppers, millipedes, mosquitoes, sod webworms (lawn moths), sowbugs (pillbugs) and ticks—Use 1 cupful in 15 to 20 gallons of water for each 500 sq. ft. of established lawn area. Use garden hose sprayer or pressure type equipment and apply full water volume to insure good penetration of turf. For best results, mow lawn and apply immediately after rain or watering. Repeat 2 to 3 weeks later if necessary.

**FRUIT INSECT CONTROL—**Recommended dosages refer to pounds per 100 gallons of water and tablespoonfuls per 1 gallon of water. Keep spray well agitated. Make full spray coverage.

**APPLES & PEARS:** San Jose Scale, Oystershell Scale, Codling Moth, Red-Banded Leaf Roller and Green Fruitworm—Apply 1 lb. per 100 gal. (2 lbs. per gal.) for codling moth. Apply 2 lbs. per 100 gal. for other insects mentioned. Repeat treatment as necessary. For optimum scale control apply when crawlers are present. Application of Sevin within 80 days after full bloom may cause fruit thinning. This may be avoided by delaying use until third cover. May cause foliar injury if used before second cover on York and McIntosh apples. The above recommendations are for areas East of the Rocky Mountains. Do not apply within 1 day of harvest.

**PLUMS, PRINES and CHERRIES:** Apply 2 lbs. per 100 or 2 lbs. per gal. water for control of Codling Moth, Forbea Scale, Brown Soft Scale, San Jose Scale, Oystershell Scale and Lecanium Scale. Apply when scale crawlers are present. For codling moth control follow the regular state fruit tree schedule. Do not apply within 1 day of harvest.

PACKAGED BY  
 SELCO SUPPLY COMPANY  
 EATON, COLORADO 80615