



6 2 71  
11 2

**DIRECTIONS FOR USE**

SELCO 0.5% DUST is suitable for direct application by any conventional dusting equipment, such as rotary or hand dusters, puff dusters, or by sprinkling from a can.

**RECOMMENDED APPLICATIONS  
SHEEP**

**Keds (Ticks), Fleeceworms (wool maggots) — HAND DUSTER** Apply 1 to 2 oz. of dust per head. Work dust well into wool over entire body

**POWER DUSTER** — Machine settings: 1½ to 4 lbs. of dust per minute. Application rate should be for treatment of 50 to 100 sheep per minute. Repeat as necessary but not within 15 days of slaughter.

**POULTRY**

**Northern fowl mite, Lice DIRECT APPLICATION DUST** birds thoroughly, especially under wings and around vent. Supplemental control can be obtained by also treating litter and housing. Repeat as necessary but not more often than weekly.

**Northern fowl mite, Lice - DUST BOX SELF-TREATMENT** Place specified dosage in shallow box for self-treatment using 2½ lbs. per 50 birds. Supplemental control can be obtained by treatment of litter and housing. Repeat as necessary.

**Northern fowl mite, Poultry red mite, Lice, Flies LITTER AND HOUSING TREATMENT** Apply 1 lb. per 70 sq. ft. of surface area to walls, ceiling, floors, roosts, nests, and adjacent areas, as well as per 20 sq. ft. of litter area. Force dust into all cracks and crevices. Repeat as necessary.

**FORMULATED BY**

**SELCO SUPPLY COMPANY • EATON & SEVERANCE, COLORADO**

**0.5% ANIMAL INSECTICIDE DUST**

**CONTAINS 0.5% CO-RAL®**

**FOR CONTROL OF ANIMAL PARASITES**

**ACTIVE INGREDIENTS:**

O, O-Diethyl O-3-chloro-4-methyl-2-oxo-2H-1-benzopyran-7-yl phosphorothioate\* ..... 0.5%

**INERT INGREDIENTS** ..... 99.5%

100.0%

\* Registered Trademark Chemagro Corporation  
U.S. Patent No. 2,748,146 — Canadian Patent No. 523,206.  
**USDA REG. NO. 1348-182**

**Caution: Keep out of reach of children!**

**CAUTION** - Avoid inhalation of dust. Wash after handling. Do not contaminate feed or food. If poisoning occurs, obtain prompt medical aid. Exposure will result in cholinesterase depression. Atropine sulfate is antidotal.

**RESTRICTIONS**

Do not treat birds within 10 days of vaccination or other stress influence. Avoid contamination of feed, water and water utensils. Provide thorough ventilation while dusting. For external insecticidal use on animals.

**NOTE:** Buyer assumes all risk of use, storage or handling of this product when not used in strict accordance with directions given herewith

**NOTICE:** Containers should be disposed of by punching holes in them and burying with wastes in non-crop lands away from water supplies.

**NET WEIGHT 50 POUNDS**