

GENERAL DIRECTIONS

This product is suitable for use in conventional power sprayers or low pressure knapsack sprayers. Just prior to use, mix the required amount of this product in sufficient water and apply in accordance with State and Federal recommendations. Do Not treat calves under 6 mos. of age. Brabma Cattle should not be treated as they may show hypersensitivity to organic phosphates.

DIRECTIONS FOR LIVESTOCK PEST CONTROL

For control of Stable Flies, Horn Flies, House Flies, Lice and Lone Star Ticks on Dairy Cattle, Beef Cattle, Sheep Goats and Hogs: Mix 1 quart of this product per 6 gals. of water (1% Concentration). Spray to thoroughly cover all parts of the animal, including the legs, using 1-2 pts. of spray per large animal and proportionately less for smaller animals.

For pen or corral operations involving spray applications to livestock with high pressure equipment, do not exceed a 0.5% concentration (1 qt. to 12 gallons of water). Spray to thoroughly cover the animal's body and legs (avoid excessive spraying of the head) using 1-2 qts. of spray per large animal and proportionately less for smaller animals. Improved Face Fly control may be achieved by adding sugar to the spray solution at the rate of 1 pound per 6-10 gallons of spray. Repeat above application as necessary to maintain control, but not more often than once every 7 days.

LOW VOLUME APPLICATION FOR CONTROL OF FACE FLIES: Mix 2 table-
spoons (1 fluid ounce) of this product per 1½ pts. of water (1% concentration). Using a calibrated sprayer, apply 5 ml. of spray to the Face and a total of 5-10 ml. to the back and sides of the animal 3 times per week. Improved control may be achieved by the addition of 1 level tablespoon of sugar (0.4 ounce) per pt. of spray solution.

FOR CONTROL OF HOUSE FLIES, FACE FLIES AND STABLE FLIES IN AND AROUND LIVESTOCK BUILDINGS: Mix 1 qt. of this product per 6 gals. of water (1% concentration). Spray inside and outside Barn Walls (including Dairy Barns), Eaves, Fences and other resting and breeding places at least twice monthly using approximately 1 gallon of finished spray per 1000 square feet. Improved House Fly and Face Fly control may be achieved by adding sugar to the spray solution at the rate of 1 pound per 6-10 gals. of spray.

NOTICE: Buyer assumes all risk of use, storage or handling of this product when not in strict accordance with directions given herewith.

NET CONTENTS 1 GALLON

DELOC

CIODRIN® INSECTICIDE

DAIRY and LIVESTOCK SPRAY

(Contains 2.0 lbs. CIODRIN Insecticide per Gal.)
For control of Horn Flies, Face Flies, Stable Flies,
and other insects listed.


EPA Reg. No. 1348-185

EPA EST. NO. 1348 — CO — 1

ACTIVE INGREDIENTS:	
Dimethyl phosphite of alpha-nedyl-	by weight
benzyl 2-hydroxy-cis-crotonate	2.000
Petroleum Hydrocarbon	1.000
INERT INGREDIENTS	27.000
TOTAL	30.000

WARNING

Keep out of reach of children!
See side panel for additional Cautions.



FOR CONTROL OF STABLE FLIES, HORN FLIES, HOUSE FLIES, AND FACE FLIES ON DAIRY and BEEF CATTLE: Mix 4 tablespoons (2 fluid ounces) of this product per 1½ pts. of water (2% concentration). Spray to thoroughly cover all parts of the animal, including the legs, using 1-2 fluid ounces of spray per animal. Apply daily with conventional hand or power sprayers. Improved control may be achieved by the addition of 1 level tablespoon of sugar (0.4 ounce) per pt. of spray solution.

FOR CONTROL OF LICE AND WINTER TICKS ON DAIRY CATTLE, BEEF CATTLE, SHEEP, GOATS AND HOGS: Mix 1-2½ pts. of this product per 3 gals. of water (0.1%-0.25% concentration). Spray animals thoroughly using up to 2 gallons of spray for the 0.1% concentration and up to 1 gallon of spray for the 0.25% concentration. Apply a second application 14 days later. Repeat applications as necessary to maintain control.

BACK-RUBBER APPLICATIONS: For use in Back-Rubbers or Face-Rubbers. For the control of HORN FLIES and FACE FLIES: Mix 1 qt. of this product per 7 gallons of Back-Rubber Base Oil (1% concentration). This may be diesel fuel or combination of Diesel fuel and non-detergent motor oil in accordance with the State Experimental Station or the State Agricultural Extension Service for additional information consistent with label directions and cautions.

PRECAUTIONS IN USING

WARNING: HAZARDOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling and before eating and smoking. Wear clean clothing. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water. For eyes, get medical attention. Keep away from heat and open flame. Atropine is antidotal.

Do not contaminate feed or foodstuffs. During commercial or prolonged exposure in spray-mixing and loading operations wear clean synthetic rubber gloves and a mask or respirator of a type found suitable by the U.S. Bureau of Mines for protection against CHLORIN insecticide. Use this material only for recommended purposes and at recommended dosages.

ENVIRONMENTAL WARNING

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. Decontaminate and destroy used containers. Do not re-use for any purpose. Decontaminate containers by rinsing thoroughly with water and alkaline detergent. Destroy containers by breaking and burying fragments in isolated areas away from crops. Dispose of rinsings in a way not to constitute a hazard or contaminate water supplies. Do not use, pour, spill or store near heat or open flame.

FORMULATED BY

**WELCO SUPPLY COMPANY
FITCH, COLORADO 80615**