

6/20/72

FEDERAL INSECTICIDE
SCIENTIFIC
NAMES REGISTER

1348-195

SELCO

TOXAPHENE 6 SPECIAL

LIVESTOCK SPRAY AND DIP

EPA Reg. No. 1348-195

ACTIVE INGREDIENTS:

Toxaphene	57.91%
Xylene	21.15%

INERT INGREDIENTS

TOTAL	100.00%
-------	---------

Technical Chlorinated Camphene (Chlorine Content 67% to 69%). Contains 6 lbs. Toxaphene per gallon.

WARNING! *Keep out of reach of children!*
See Side Panel for Additional Directions!
and Warnings!

NET CONTENTS 1 GALLON

6/20/72

U.S. DEPARTMENT OF AGRICULTURE
BUREAU OF PLANT INDUSTRY
OFFICE OF ENTOMOLOGICAL RESEARCH
REGISTERED UNDER NO. 1348-195

A Special Formulation for the control of SARCOPTIC MANGE on Cattle and PSOROPTIC SCAB on Cattle and Sheep. Other uses listed below.

RECOMMENDED USES

RANGE and FEEDER CATTLE: For control of Sarcoptic Mange, Chorioptic Mange, Psoroptic Scab, Cattle Lice, Ear Tick, Lone Star Tick, Cattle Fever Tick, Winter Tick and Horn Flies - Use 1 gallon of this product to 150 gallons of water and stir or agitate well to form a creamy emulsion. Spray or dip animals thoroughly to assure penetration to the skin. For spraying operation it is advisable to use a high pressure sprayer at a minimum of 250 pounds pressure with constant agitation of spray mixture in the tank. If spray-dip machines are used, we recommend that the manufacturers recommendations and directions for operation be used. It is essential that suction strainers be removed and cleaned as needed to remove hair and debris to permit recycling of the spray mixture.

For use in dipping vats, the same concentration is used as given above. Before dipping operation, thoroughly mix the vat contents by using a pole or bucket attached to a rope. Addition of a wetting agent is advisable if the water is known to be extremely hard and highly alkaline. Standard methods used for determining dipping vat concentration of toxaphene emulsions may be used for this product. Animals in scabies infected herds or flocks are dipped twice; those in exposed herds or flocks at least once. The interval between the first and second dipping is 10-14 days. NOTE: Do not treat animals under three months of age. Do not dip animals under 6 months of age. Avoid treating sick or overheated animals. Avoid spray or dip treatment during stormy or cold weather. If treating in warm weather, provide shade for animals after treatment. Water animals before dipping or spraying to prevent them from drinking the dip or puddles from the spray. In case of quarantine where calves require treatment, spray very lightly. Do not treat animals within 28 days of slaughter.

BACK RUBBERS: Beef Cattle - As an aid for the control of Lice and Horn Flies, mix 1 gallon to 15 gallons of fuel oil or mixture of mineral oil and soak sack or cloth of burlap rubbing device or fill compartment of back rubber as required. Recharge back rubber as necessary. Mount or place back rubber where it is readily accessible to animals as they move to feed, water or pass through gates from one pen or pasture to another. CAUTION: Do not use on calves under six months of age or around dairy animals. Do not use back rubber mixture as a spray. Do not permit animals access to treatment with back rubbers within 28 days of slaughter.

SHEEP and GOATS: For control of Psoroptic Sheep Scab - Use at the rate of 1 gallon to 150 gals. of water. Thoroughly wet animals. For control of Sheep Ticks (Keds), Sheep Lice and Goat Lice - Use 1 gallon to 300 gallons of water for dip and 1 gallon to 150 gallons of water for a spray. Wet animals thoroughly. Do not treat sick animals. Do not treat milk goats. Do not use on sheep or goats within 28 days of slaughter.

WARNING

May be fatal if swallowed! Do not breath spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Do not apply in dairy barns or on dairy animals. Do not apply in dairy barns or on dairy animals. Do not use on poultry or in poultry houses.

TOXIC TO FISH AND WILDLIFE. Keep out of lakes, ponds and streams! Equipment and Containers: Rinse equipment and containers and dispose of wastes by burying in non-crop lands, away from water supplies. Containers should be disposed of by punching holes in them and crushing and burying with wastes or burning. (Keep out of Smoke.)

NOTICE: Buyer assumes all risk of use, storage or handling of this product when not in strict accordance with directions given hereon.

Do not use, pour, spill or store near heat or open flame.

Formulated By
SELCO SUPPLY COMPANY
EATON, COLORADO 80615