

GENERAL DIRECTIONS

Apply this product when insects or their damage appear. Make application in the morning or evening when it is cool. Treat upper and lower surface of all foliage for complete coverage. Repeat as necessary, generally at 7-day intervals. For best results, apply with a dust gun or garden duster. Do not apply in excess.

VEGETABLE INSECT CONTROL

BROCCOLI, BRUSSEL SPROUTS, CAULIFLOWER, and HEAD LETTUCE: To control flea beetles, harlequin bug, armyworms and Corn Earworm. Do not apply within 3 days of harvest.

BEANS (green beans, lima beans, snap beans): To control Mexican bean beetle, cucumber beetles, flea beetles. No post-treatment limitations on application to beans.

CORN (Sweet): To control Corn Earworm attacking silks and ears, apply at 3 to 5 day intervals starting when first silks appear, and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestations. Timing and good coverage are essential for effective control. No time limitation to harvest. Note: Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

CUCUMBERS, MELONS, PUMPKIN, SQUASH: To control pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers and squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use on watermelons in Florida. No time limitation between application and harvest.

POTATO, TOMATO, EGGPLANT, PEPPER: To control Colorado Potato Beetle, Flea Beetles, Leafhoppers, Tomato Hornworm, Plant Bugs. No time limitation between application and harvest.

STRAWBERRIES: To control strawberry leaf roller and strawberry weevil. Do not apply within 1 day of harvest.

ORNAMENTAL INSECT CONTROL — This product can be used on a wide range of ornamental plants, flowers, shrubs and small trees — rose, carnation, gladiolous, chrysanthemum, lily, hydrangea, dogwood. Do not use on Boston Ivy. Apply to the upper and lower leaf surface of all foliage for complete coverage to control — Bagworm, Boxelder bug, Elm leaf beetle, Fleabeetles, Leaf roller, Mealy bugs, Periodical cicada, Plant bugs, Scale insects, Tent caterpillars, Willow leaf beetles. NOTE: Sevin does not control Spider mites. Where infestations are encountered, use a miticide as recommended. For protection of honeybees avoid use if possible during periods when honeybees are visiting the crop. Injury to tender foliage may occur if applications are made when plants are wet or in the presence of high humidity of several days duration following treatment.

Formulated by
SELCO SUPPLY COMPANY
EATON, COLORADO 80615

SELCC

SEVIN 5% DUST

A MULTIPLE USE INSECTICIDE

For The Control of Certain Insects on Vegetables, Berries
and Ornamentals

ACTIVE INGREDIENT:	
Carbaryl (1-naphthyl N - methylcarbamate)	5.0%
INERT INGREDIENT	95.0%
TOTAL	100.0%

*Registered trade mark of Union Carbide Corporation

EPA REG. NO. 1348-204

CAUTION: KEEP OUT OF REACH OF CHILDREN
See Back Panel for Additional Cautions and Directions.

HARMFUL IF INHALED OR SWALLOWED.

Do not Breathe Dust. Do not Take Internally. Avoid Skin Contact. Wash thoroughly after using, and before eating. Avoid Contamination of Food and Food-stuffs, Feed troughs and water receptacles. NOTE TO PHYSICIAN: Carbaryl is a moderate reversible, Cholinesterase inhibitor. Atropine is antidotal.

NOTE: Do not reuse empty container. Rinse with water and discard by burying in non-crop land away from water supply.

NOTICE: Buyer assumes all risk of use, storage or handling of this product when not in strict accordance with directions given herewith.

HOME AND GARDEN PACKAGE

NET WEIGHT 1 POUND

ACCEPTED

1348-204