

CLOROX

Mc SEED PROTECTANT

ACTIVE INGREDIENT

Trichlorophenyl Ether

- 100% Trichlorophenyl Ether
- 100% Trichlorophenyl Ether
- 100% Trichlorophenyl Ether

INERT INGREDIENTS

Water
Alcohol
Trichlorophenyl Ether

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Inhaled or absorbed on skin. Avoid inhalation of vapor or fumes. Avoid contact with skin, eyes or clothing. In case of contact with skin or eyes, flush with plenty of water. If you get medical attention, wash with water. After feeding, and before putting on clothes, wash your hands. Avoid contact with children and pets.


This product is not for use. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Contents 1 U.S. Quart

CLOROX SEED PROTECTANT

CLOROX SEED PROTECTANT

CLOROX SEED PROTECTANT



EVERSHIELD Mc Seed Protectant is primarily designed to aid in protection of stored seed by reducing losses from the following insects: cadelle, rice weevil, confused flour beetle, Angoumois grain moth and Indian meal moth.



Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

- For Use By Professional:
1. Add EVERSHIELD to cool tap water; prepare the EVERSHIELD mixture.
 2. Add 1.25 oz. Dye to each dilution and mix.
 3. Add mixture and dye to seed applicator as directed.

U.S. PATENT NO. 3,113,399

Mc SEED PROTECTANT

E.P.A. REG. NO. 1352-16

IMPORTANT

Treated seed must not be used for human consumption or as animal feed. Bags containing treated seed should be labeled.

TREATED SEED - DO NOT USE FOR FEED, FOOD OR OIL PURPOSES

Some States and the Federal Government have regulations for the application of special tags or labels to be attached to bags containing treated seed.

Contents 1 U.S. Quart

ACTIVE INGREDIENT

*Methoxychlor, Technical 38.0%

*Equivalent to 33.44% 2,2-bis (p-methoxyphenyl) -1,1,1-trichloroethane and 4.56% other isomers and related compounds.

INERT INGREDIENTS: 62.0%

Pounds of Active Ingredient Per Gallon of EVERSHIELD Mc SEED PROTECTANT

*Methoxychlor, Technical 3.75 lb/gal.

CARGILL, INCORPORATED
Cargill Building
Minneapolis, Minnesota 55402

CAUTION!

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through skin. Avoid inhalation of vapor or fumes. Avoid contact with skin, eyes or clothing. In case of contact with skin or eyes, flush with plenty of water, for eyes get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Avoid contamination of feed and food-stuffs.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

GENERAL INFORMATION

Evershield is dustfree, binds protectant firmly to seed, in a water permeable film, improves seed appearance, eliminates buildup in planter plates, dries fast without need of additional heat.

Registered trademark of Cargill Inc.

**STORE IN COOL DRY PLACE
DO NOT REUSE CONTAINER;
DESTROY WHEN EMPTY
DO NOT FREEZE**

Seed

Beans (Snap & Dry), Soybeans, Cowpeas, Corn (Field), Peas, Bean (Lima), Sesame, Buckwheat, Sorghum, Millet, Cantaloupe, Cucumber, Pumpkin, Squash, Watermelon, Castor Bean, Cotton, Acid delinted, Wheat, Rye, Corn (Sweet), Eggplant, Okra, Tomato, Barley, Flax, Beet (Garden), Broccoli, Brussels Sprouts, Cabbage, Carrot, Cauliflower, Grasses, Kale, Kohlrabi, Endive, Lettuce, Pepper, Radish, Spinach, Swiss Chard, Turnip, etc.

For Further Information, See Chapter 1, Section 1.1, of the Federal Register, Vol. 32, No. 1, dated January 1, 1967, and subsequent amendments.

ACCEPTED

Aug - 13 - 1977

1352-16

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROVEMENTAL ACT
NOTICE REGISTERED
UNDER NO.

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.



SEED PROTECTANT

E.P.A. REG. NO. 1352-16

CAUTION!

**KEEP OUT OF REACH
OF CHILDREN**

Harmful if swallowed, inhaled or absorbed through skin. Avoid inhalation of vapor or fumes. Avoid contact with skin, eyes or clothing. In case of contact with skin or eyes, flush with plenty of water, for eyes get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Avoid contamination of feed and food-stuffs.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

GENERAL INFORMATION

Evershield is dustfree, binds protectant firmly to seed, in a water permeable film, improves seed appearance, eliminates buildup in planter plates, dries fast without need of additional heat.

Registered trademark of Cargill Inc.

**STORE IN COOL DRY PLACE
DO NOT REUSE CONTAINER;
DESTROY WHEN EMPTY
DO NOT FREEZE**

For Use By Professional Seed Treaters Only

- 1 Add EVERSHIELD Mc Seed Protectant concentrate to cool tap water and mix thoroughly for 15 minutes to prepare the EVERSHIELD Mc dilution.
- 2 Add 1.25 oz. of EVERSHIELD Seed Protectant Liquid Dye to each 5 gallons of the EVERSHIELD Mc dilution and mix for an additional 10 minutes.
- 3 Add mixture of EVERSHIELD Seed Protectant, water and dye to seed at the following rates in any seed applicator assuring complete and uniform coverage.

Seed	Fluid ozs. of Conc. Diluted to 1 Gallon	Application Rates (Cubic Centimeters of Diluted Product per 100 lbs. of Seed)
Beans (Snap & Dry)	9	240
Soybeans, Cowpeas	11	270
Corn (Field)	9	370
Peas, Bean (Lima)	10	320
Sesame	75	480
Rice	10	420
Sorghum, Millet	12	420
Cantaloupe, Cucumber, Pumpkin,	1.95	250
Squash, Watermelon	12	420
Castor Bean	18	270
Cotton - Acid delinted	14	370
Wheat	12	480
Rye	1.55	420
Corn (Sweet)	19	320
Eggplant, Okra, Tomato	14	530
Barley		
Flax		
Beet (Garden), Broccoli, Brussel Sprouts,		
Cabbage, Carrot, Cauliflower, Collards,		
Grasses, Kale, Kohlrabi, Endive, Mustard,		
Lettuce, Pepper, Radish, Small seed		
legumes, Spinach, Swiss Chard, Turnip		

21CFR Chapter 1 Section 3.13 requires grain seed treated with poisonous substances in excess of recognized tolerances be suitably colored to prevent their subsequent inadvertent use as food for man or feed for animals.

IT
nical 38.0%
% 2,2-bis
1,1,
d 4.56%
lated
FS: 62.0%
edient
HIELD
ANT
nical 3.75 lb/gal.
ORPORATED
ding
Minnesota
55402