NOTICE:

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combinations with other materials.

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. FLO PRO Seed Protectant Concentrate is for use by professional seed treaters only. Observe all cautions and limitations on labeling of products used in mixtures.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment. Do not reuse empty container. Open dumping is prohibited.

PESTICIDE DISPOSAL

Pesticide, spray mixture or rinsate that cannot be used or chemical reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL
Triple rinse (or equivalent) and dispose in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

Consult federal, state or local disposal authorities for approved, alternative procedures such as limited open burning.

STORAGE

Store in a cool dry place.

Contents: U.S. 5 Gallons

APR 1 0 1981

Under the Federal Insecticide.
Fungicide, and Rodenticide Act.
as amended, for the pesticide
Togistered under
TPA Reg. No.

ACTIVE INGREDIENTS:

Captan* 28.37% Related derivatives......1.63% INERT INGREDIENTS70.0%

*N-Trichloromethylthio-4-cyclohexene-1, 2-dicarboximide

Pounds of Active Ingredient Per Gallon of FLO-PRO C Seed Protectant.

Captan2.94 lbs./gal.



Manufactured by Chemical Products Division Agricultural Products Dept. Cargill, incorporated P.O. Box 9300 Minneapolis, Minnesota 55440

EPA REG NO. 1352-18 EPA EST. NO. 36628-TN-1 DANGER

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS STATEMENT OF PRACTICAL TREATMENT

If swallowed: Harmful if swallowed, get medical

If inhaled: May be harmful if inhaled, get medical attention. Avoid inhalation by using a attention.

Skin: Avoid conte Swith skin or clothing. Wear rubber gloves. Wash affected area thoroughly with soap and water. Wash hands and face thoroughly with soap and water after use and before eating or smoking. respirator.

Eyes: Avoid contact with eyes. Wear protective eye covering. In case of contact with eyes, this immediately with running water for 15 minutes and get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS
This product is toxic to tish. Keep out of takes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Flo-Pro is a Trademark of Cargill, Inc.

Flo-Pro C Seed Protectant is for use by professional seed treaters only.



..... 1.63%

tnio-4-cyclohexene-1,

redient Per Gallon of

at Products Division

tural Products Dept. . Incorporated

apolis, Minnesota 55440

A REG NO. 1352·18

EST. NO. 36628-TN-1

.....2.94 ibs./gal.

TS:

⊬e**ctant.**

actured by

APR 1 0 1981

Under the Federal Insacticide. Fungicide, and Rodenticide Act. cs amended, for the pesticide EPA Reg. Ilo. 1352-18

DANGER 28.37%

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS STATEMENT OF PRACTICAL TREATMENT

If swallowed: Harmful if swallowed, get medical

attention.

If inhaled: May be harmful if inhaled, get medical attention. Avoid inhalation by using a respirator.

Skin: Avoid contact with skin or clothing. Wear skin: Avoid contact with skin or clothing. Wear rubber gloves. Wash affected area thoroughly with soap and water. Wash hands and face thoroughly with soap and water after use and before eating or smoking.

Eyes: Avoid contact with eyes. Wear protective eye covering. In case of contact with eyes, flush immediately with running water for 15 minutes and get medical attention if irritation persists.

persists.

ENVIRONMENTAL HAZARDS
This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Flo-Pro is a Trademark of Cargill, Inc.

d Protectant is for use by professional ser 1 treaters only.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

. Treated seed must not be used for human consumption or as animal feed. Bags containing treated seed should be labeled: TREATED SEED - DO NOT USE FOR FEED, FOOD OR OIL PURPOSES.

Primarily designed for the protection of seed from soil borne fungi causing seed, seedling and root rots.

Store in a cool dry place.

Apply with commercial seed treating equipment such as a slurry treater.

1. Add 1 gallon of FLO-PRO Seed Protectant concentrate to 5 gallons of cool tap water and mix thoroughly for 15 minutes to prepare the FLO-PRO dilution.

2. Add 7 oz. of CARGILL Seed Protectant Liquid Dye** to each 6 gallons of the FLO-PRO dilution and mix for an additional 10 minutes.

Seed

Application Rates (Cubic Centimeters of Diluted Product Per 100 lbs. of Seed)

Sweettsew2
Wheat*270
Oats*420
Barley*320
Flax530
Peas370
Cucumber, Cantaloupe, Tomato 420
Squash, Pumpkin, Watermelon, Musk-
melon280
Cotton Acid Delinted 420
Reginned
Cabbage, Cauliflower, Broccoli, Brussels
Sprouts, Collards, Kale, Mustard,
Turnip
Peppers (Calif. Wonder)420
Cowpeas420
Rice
Rye*370
Crucifers210
*For Seed & Seedling Rots, not for Bunt

Smut.

1. Add 1 gallon of FLO-PRO Seed Protectant concentrate to 3 gallons of cool tap water and mix thoroughly for 15 minutes to prepare the FLO-PRO dilution.

2. Add 7 oz. of CARGILL Seed Protectant Liquid Dye** to each 4 gallons of the FLO-PRO dilution and mix for an additional to dilution and the second s tional 10 minutes.

Seed

Application Rates (Cubic Centimeters of Diluted Product Per 100 lbs. of Seed)

Beans (except baby limas)	50 50
Sorghum - hulled (Kernel Smut) 42 Sugar Beets - West	20
Spinach5	7Ö

1. Add 1 gallon of FLO-PRO Seed Protectant concentrate to 2 gallons of cool tap water and mix thoroughly for 15 minutes to prepare the FLO-PRO dilution.

2. Add 7 oz. of CARGILL Seed Protectant Liquid Dye** to each 3 gallons of the FLO-PRO dilution and mix for an additional 10 minutes.

tional 10 minutes.

Seed

Application Rates (Cubic Centimeters of Diluted Product Per 100 ibs. of Seed)

Clover, Alfalfa, Lespedeza, Trefoil Beets, Swiss Chard, Sugar Beets-	. 580
East	. 850
Blue Grass, Grasses	. 580

1. Use Undiluted Concentrate and add 1 oz. CARGILL Seed Protectant Liquid Dye** to each gallon of the FLO-PRO Concentrate. Mix for an additional 20 minutes.

Seed

Application Rates (Cubic Centimeters of Product Per 100 lbs. of Seed)

**21CFR Chapter 1 Section 3.13 requires grat's seed treated with poisonous substances in excess of recognized tolerances be suitably colored to prevent their subsequent inadvertent use as food for man or feed for animals.