TU-
CNB
TM

protectant

INGREDIENT:

|	
chloronitrobenzene	24.0%
INGREDIENTS:	76.0%
	100.0%

s 2.23 lbs. of Pentachloronitrobenzene per llon

OUT OF REACH OF CHILDREN

4   DANGER

for additional precautionary statements.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Using suitable liquid or slurry seed treating equipment such as Panogen or Mist-O-Matic, apply RTU-PCNB at rates shown below.

Seed should be sound, well cured and cleaned prior to treatment. Use only the recommended rates of product per 100 pounds of seed.

Planting Seeds	Application Rates Per 100 Pounds of Seed	Seedling Disease Complex
BARLEY	3.9-7.8 fl. oz. (115-231 cc)	Covered Smut
OATS	5.8-11.6 fl. oz. (172-343 cc)	Oat Smut
WHEAT	3 fl. oz. (89 cc)	Common Smut or Bunt and Damping-off caused by Rhizoctonia
COTTON	Acid Delinted 10.8 fl. oz. (320 cc) Mechanically Delinted 14.7 fl. oz. (435 cc)	Seed rot and early stages of Pre-emergence damping-off caused by Rhizoctonia

At the recommended rates, apply RTU-PCNB undiluted directly from the drum to the seed. If desired, add one part water to five parts RTU-PCNB and recalibrate equipment so that recommended rates will still be applied. Slight agitation may be required to mix the material. Keep container tightly closed when not in use. Do not feed treated seed to animals.

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food, or feed by storage or disposal.
Container Disposal: Rinse containers and dispose of in an incinerator or landfill approved for pesticide containers.
General: Consult Federal, State or local disposal authorities for approved alternative procedures such as limited open burning.

NOTICE TO BUYER

The product conforms to its chemical description and is fit for the use intended when used in accordance with label directions. Cargill will not be responsible for special, consequential or contingent damages arising from injury to persons or property.